

RATIONAL SUBSETS IN HNN-EXTENSIONS AND AMALGAMATED PRODUCTS

Markus Lohrey

*Universität Stuttgart, FMI, Germany
lohrey@informatik.uni-stuttgart.de*

Géraud Sénizergues

*LABRI, Université Bordeaux I, France
ges@labri.fr*

Several transfer results for rational subsets and finitely generated subgroups of HNN-extensions $G = \langle H, t; t^{-1}at = \varphi(a)(a \in A) \rangle$ and amalgamated free products $G = H *_A J$ such that the associated subgroup A is finite. These transfer results allow to transfer decidability properties or structural properties from the subgroup H (resp. the subgroups H and J) to the group G .

1. Introduction

This paper is the first paper in a series of papers dealing with algorithmic problems for equations over HNN-extensions [HNN49] and amalgamated free products of groups. These two operations are of fundamental importance in combinatorial group theory [LS77]. Recall that an amalgamated free product

$$G = \langle H_1 * H_2, a = \varphi(a)(a \in A) \rangle \quad (1)$$

of two groups H_1 and H_2 with subgroups $A \leq H_1$, $B \leq H_2$ and an isomorphism $\varphi : A \rightarrow B$, results from the free product $H_1 * H_2$ by identifying every element $a \in A_1$ with $\varphi(a) \in A_2$. We also say that A and B in (1) are the identified subgroups. An HNN extension

$$G = \langle H, t; t^{-1}at = \varphi(a)(a \in A) \rangle \quad (2)$$

of a group H , where A and B are isomorphic subgroups of H and $\varphi : A \rightarrow B$ is an isomorphism, results from adding to H a new generator t such that the conjugation of the subgroup $A \leq H$ by t realizes the isomorphism φ . We also say that A and B in (2) are the associated subgroups.

One of the first important applications of HNN-extensions was a more transparent proof of the celebrated result of Novikov and Boone on the existence of a finitely presented group with an undecidable word problem, see e.g. [LS77]. Such a group can be constructed by a series of HNN-extensions starting from a free group. This shows that arbitrary HNN-extensions do not have good algorithmic properties. In

this paper, we restrict to HNN-extensions (resp. amalgamated free products), with finite associated (resp. identified) subgroups. This is an important special case. For instance, a famous result of Stallings states that a group G has more than one end if and only if it is either an HNN-extension with finite associated subgroups or an amalgamated free product with finite identified subgroups [Sta71]. Moreover, a group is virtually-free (i.e., has a free subgroup of finite index) if and only if it can be built up from finite groups using amalgamated free products with finite identified subgroups and HNN-extensions with finite associated subgroups [DD90]. Finally, many important structural and algorithmic properties of groups are preserved by HNN-extensions (resp. amalgamated free products) with finite associated (resp. identified) subgroups, see e.g. [AG73,Bez98,KSS06,KS70,KS71,KWM05,MR04].

In this paper, we will prove several such preservation results for rational subsets and finitely generated subgroups of HNN-extensions and amalgamated free products of groups. Recall that the set of rational subsets of a monoid \mathbb{M} is the smallest class of subsets of \mathbb{M} which contains all finite subsets of \mathbb{M} and which is closed under union, product and the Kleene star (which associates with a subset $U \subseteq \mathbb{M}$ the submonoid of \mathbb{M} generated by U). By Kleene's theorem, the rational subsets of a free monoid are a Boolean algebra. This result also holds for free groups [Ben69] and even virtually-free groups [Sén96]. Further results on rational subsets of groups can be found in [Gru90,KSS06,Ned00].

Some of our results are already known, others appear to be new. Apart from proving these new results, the purpose of our investigations is two-fold: first, we prepare the background for our further considerations on equations over HNN-extensions and amalgamated free products [LS06a,LS06b]. Second, we present a unified automata-theoretic framework, which is partly inspired by [Reu85,Sak87,Sak92], for reasoning about rational subsets of HNN-extensions and amalgamated free products. This framework allows us to deduce transfer theorems in a unified way. Moreover, our approach allows us to prove results about rational subsets even for HNN-extensions and amalgamated free products of *monoids*, in case the associated parts A and B in (1) and (2) are groups. Let us now describe in more detail the content of this paper.

Section 2 contains necessary preliminaries concerning monoids, groups, and rational sets. Section 3 contains basic definitions and facts for HNN-extensions. In Section 4.1, we introduce finite t -automata, which are a specific automaton model in order to recognize subsets of an HNN-extension $G = \langle H, t; t^{-1}at = \varphi(a)(a \in A) \rangle$, where H is a monoid and A and $B = \varphi(A)$ are finite groups. The transitions of such an automaton are either labelled with t , t^{-1} , or sets from some subset $\mathcal{F} \subseteq \mathcal{P}(H)$ (which has to contain every singleton subset consisting of an element from $A \cup B$). For our specific applications concerning rational subsets (resp. finitely generated subgroups) of the HNN-extension G , \mathcal{F} is the set of rational subsets of H (resp. the set of left cosets of finitely generated subgroups of H). In Section 4.2–4.3 we prove several basic facts about finite t -automata and in Section 4.4 we prove two

fundamental normalization results for finite t -automata (Propositions 22, 28 and 33). Roughly speaking, these normalization results state that we can always enforce some nice properties for finite t -automata. Based on these results we can show in Section 4.5 Theorem 37, which is our main result for rational subsets of HNN-extensions. It says that the following four properties can be transferred from the monoid H to the HNN-extension $G = \langle H, t; t^{-1}at = \varphi(a)(a \in A) \rangle$ (with A and B finite groups):

- (a) The set of rational subsets is closed under intersection.
- (b) The set of rational subsets is closed under complement (i.e., it is a Boolean algebra).
- (c) The membership problem for rational subsets is decidable.
- (d) The emptiness problem for boolean combinations of rational subsets is decidable.

Statement (c) is covered by [KSS06, Theorem 5.1] in the case that H (and hence G) is a group, whereas (a), (b), and (d) are new to the knowledge of the authors. It should be remarked that properties (a) and (b) (resp. (c) and (d)) are independent in the following sense: Clearly, closure under complement for the rational subsets implies closure under intersection. On the other hand, in Appendix B, we construct a finitely generated monoid for which the rational subsets are closed under intersection but are not closed under complement (we don't know, whether a finitely generated group with this property exists as well). Similarly, decidability of the emptiness problem for boolean combinations of rational subsets implies decidability of the membership problem for rational subsets. On the other hand, for the group $F \times \mathbb{Z}$ (where F is the free group of rank 2), the membership problem for rational subsets is decidable [KSS06], whereas the inclusion problem for rational subsets (which is a special instance of the emptiness problem for boolean combinations of rational subsets) is undecidable [AH89].

Let us mention that more general notions of HNN-extensions for monoids were defined in the literature (see for instance [How63]) where A and B in (2) are not necessarily groups. It is not clear whether our results hold for these HNN-extensions as well.

In Section 4.6 we deal with finitely generated subgroups of HNN-extensions. Here we assume that H (and hence G) in (2) is a group. Our main result is Theorem 43, which states that the following three properties can be transferred from a group H to the HNN-extension $G = \langle H, t; t^{-1}at = \varphi(a)(a \in A) \rangle$ (with A and B finite):

- (a) Decidability of the generalized word problem (i.e., the question whether a given group element belongs to a given finitely generated subgroup).
- (b) The Howson property.
- (c) The LERF property.

Recall that a group G has the Howson property if the intersection of two finitely generated subgroups is again finitely generated. The group G has the LERF property [Hal49] if for every finitely generated subgroup $H < G$ and every $g \in G \setminus H$ there exists a subgroup Z of finite index in G with $H \leq Z < G$ and $g \in G \setminus Z$. Statement (a) is covered by [KWM05, Corollary 5.15], statement (b) was shown in [Bez98,KS71], and statement (c) was shown in [MR04].

Finally, in Section 6 we briefly sketch the proof of analogous transfer results for amalgamated free products (Theorem 51 and 53). Since the treatment of amalgamated free products exactly follows our treatment of HNN-extensions, we will just state the necessary adaptations but omit most of the proof details. For the special case of an amalgamated free product of two finite groups, algorithmic problems for finitely generated subgroups were recently studied in [ME07b,ME07a] by extending Stallings's folding technique [Sta83].

2. Preliminaries

We recall in this section all needed definitions and classical results concerning groups, monoids, finite automata, and rational subsets of a monoid. We also recall some classical algorithmic problems about rational subsets of finitely generated monoids.

We assume that the reader has some familiarity with monoid presentations and group theory. For a monoid homomorphism $\pi : \mathbb{M}_1 \rightarrow \mathbb{M}_2$ we denote with $\text{Ker}(\pi) = \{(x, y) \mid \pi(x) = \pi(y)\}$ the *kernel* of π , it is a monoid congruence on \mathbb{M}_1 . The neutral element 1 of a monoid \mathbb{M} will be identified with the submonoid $\{1\}$ in the rest of this paper.

The powerset of a set S is denoted by $\mathcal{P}(S)$. For an equivalence relation \equiv on S and $s \in S$ we denote with $[s]_{\equiv}$ the equivalence class of \equiv containing s . For a subset $T \subseteq S$ let $[T]_{\equiv} = \bigcup_{s \in T} [s]_{\equiv}$. The boolean closure of a set $F \subseteq \mathcal{P}(S)$ is denoted by $\text{Bool}(F)$. It contains all subsets of S , which can be obtained by finitely many applications of complement and binary union, starting with subsets from F .

A *semi-Thue system* ST over the (not necessarily finite) alphabet X is just a (possibly infinite) subset of $X^* \times X^*$. We associate with ST the one-step rewrite relation \rightarrow_{ST} , where $x \rightarrow_{\text{ST}} y$ for $x, y \in X^*$ if and only if there are $u, v \in X^*$ and $(\ell, r) \in \text{ST}$ such that $x = ulv$ and $y = urv$. The set of irreducible words with respect to ST is

$$\text{Irr}(\text{ST}) = \{x \in X^* \mid \neg \exists y \in X^* : x \rightarrow_{\text{ST}} y\}.$$

A binary relation \rightarrow over a set E is *confluent* if for all $e, f, f' \in E$:

$$(e \rightarrow^* f \wedge e \rightarrow^* f') \Rightarrow (\exists g \in E : f \rightarrow^* g \wedge f' \rightarrow^* g)$$

Suppose that a semi-Thue system ST has no overlapping left-hand sides, i.e., for all $(\ell, r), (\ell', r') \in \text{ST}$ and $u, v \in X^*$ we have

$$(u\ell = \ell'v) \Rightarrow |u| \geq |\ell'| \quad \text{and} \quad (u\ell v = \ell') \Rightarrow (u = v = \varepsilon \wedge r = r').$$

Then it can be easily shown that the relation $\rightarrow_{\mathcal{S}\mathcal{T}}$ is confluent. See [DJ91] for more information about confluence and other properties of rewriting systems.

2.1. Finite automata over a free monoid

Let X be an alphabet (that we do not assume to be finite, in general). A finite automaton over the free monoid X^* is a 5-tuple

$$\mathcal{A} = \langle \mathcal{X}, \mathcal{Q}, \delta, \mathcal{I}, \mathcal{T} \rangle, \quad (3)$$

where \mathcal{X} is a finite subset of X , \mathcal{Q} is the finite set of states, $\mathcal{I} \subseteq \mathcal{Q}$ is the set of initial states, $\mathcal{T} \subseteq \mathcal{Q}$ is the set of terminal states, and δ , the set of transitions, is a subset of $\mathcal{Q} \times \mathcal{X} \times \mathcal{Q}$. Let

$$\text{BR}(\mathcal{Q}) = \mathcal{P}(\mathcal{Q} \times \mathcal{Q})$$

be the monoid of binary relations over the set \mathcal{Q} . We recall that, given two binary relations $m, m' \in \text{BR}(\mathcal{Q})$, their composition is defined by

$$m \circ m' = \{(q, q'') \in \mathcal{Q} \times \mathcal{Q} \mid \exists q' \in \mathcal{Q} : (q, q') \in m \wedge (q', q'') \in m'\}.$$

The automaton (3) induces a representation map

$$\mu_{\mathcal{A}} : (X^*, \cdot, \varepsilon) \rightarrow (\text{BR}(\mathcal{Q}), \circ, \text{Id}_{\mathcal{Q}})$$

defined as the unique monoid homomorphism such that for every $x \in X$:

$$\mu_{\mathcal{A}}(x) = \{(q, r) \in \mathcal{Q} \times \mathcal{Q} \mid (q, x, r) \in \delta\}.$$

We denote by \odot the natural right-action of the monoid $\text{BR}(\mathcal{Q})$ over $\mathcal{P}(\mathcal{Q})$, i.e., for every $P \subseteq \mathcal{Q}$ and $m \in \text{BR}(\mathcal{Q})$ let

$$P \odot m = \{q \in \mathcal{Q} \mid \exists p \in P : (p, q) \in m\}. \quad (4)$$

The language recognized by \mathcal{A} is

$$\text{L}(\mathcal{A}) = \{u \in X^* \mid \mu_{\mathcal{A}}(u) \cap (\mathcal{I} \times \mathcal{T}) \neq \emptyset\}.$$

A subset $L \subseteq X^*$ is called *regular* if and only if there exists some finite automaton \mathcal{A} such that $L = \text{L}(\mathcal{A})$. Given a subset $Y \subseteq X$ and two states $p, q \in \mathcal{Q}$, the automaton $\mathcal{A}[Y, p, q]$ over Y^* is defined by

$$\mathcal{A}[Y, p, q] = \langle \mathcal{X} \cap Y, \mathcal{Q}, \delta \cap (\mathcal{Q} \times Y \times \mathcal{Q}), \{p\}, \{q\} \rangle. \quad (5)$$

2.2. Rational subsets of a monoid

Let \mathbb{M} be a monoid together with a generating set X . Thus $\mathbb{M} = X^*/\equiv$, for some monoid congruence \equiv over X^* . We denote by

$$\pi : X^* \rightarrow X^*/\equiv$$

the canonical homomorphism with $\pi(u) = [u]_{\equiv}$ for $u \in X^*$.

The set

$$\text{Rat}(\mathbb{M}) \subseteq \mathcal{P}(M)$$

is the smallest subset of $\mathcal{P}(M)$ which contains the finite subsets of \mathbb{M} and which is closed under the operations \cup (the union operation), \cdot (the product operation) and $*$ (the star operation, associating with a subset $P \subseteq \mathbb{M}$ the smallest submonoid of \mathbb{M} containing P).

One can derive from Kleene's theorem about rational subsets of X^* that the rational subsets of \mathbb{M} are exactly the subsets of the form $\pi(L(\mathcal{A}))$, where \mathcal{A} is a finite automaton over the free monoid X^* . For such an automaton $\mathcal{A} = \langle \mathcal{X}, \mathcal{Q}, \delta, l, T \rangle$ let the representation map

$$\mu_{\mathcal{A}, \mathbb{M}} : \mathbb{M} \rightarrow (\text{BR}(\mathcal{Q}), \circ, \text{Id}_{\mathcal{Q}})$$

by defined by

$$\mu_{\mathcal{A}, \mathbb{M}}(m) = \bigcup \{ \mu_{\mathcal{A}}(u) \mid u \in X^*, \pi(u) = m \}$$

for $m \in \mathbb{M}$. Notice that

$$\text{Id}_{\mathcal{Q}} \subseteq \mu_{\mathcal{A}, \mathbb{M}}(1) \text{ and } \mu_{\mathcal{A}, \mathbb{M}}(m) \circ \mu_{\mathcal{A}, \mathbb{M}}(m') \subseteq \mu_{\mathcal{A}, \mathbb{M}}(m \cdot m') \quad (6)$$

but in general the map $\mu_{\mathcal{A}, \mathbb{M}}$ is *not* a monoid homomorphism. In Section 4.1, we introduce in the particular case where \mathbb{M} is an HNN-extension a new kind of finite automaton, called *finite t -automata*, which do not even fulfil property (6). We will show that, nevertheless, we can always normalize finite t -automata in such a way that the map μ fulfils some nice algebraic property, see Definition 9 and Proposition 33.

Anissimov and Seifert have shown:

Theorem 1 ([AS75]). *Let G be a group. The rational subgroups of G are exactly the finitely generated subgroups of G .*

We present a proof sketch for Theorem 1, because the underlying idea will be used later in a slightly modified form.

Proof sketch for Theorem 1. Let $G = X^*/\equiv$ be a group, where X is some alphabet and \equiv is a congruence over X^* . We suppose that X is endowed with an involution $x \mapsto x^{-1}$ such that for every $x \in X$, $xx^{-1} \equiv x^{-1}x \equiv \varepsilon$. We extend this involution to X^* by setting $(x_1 \cdots x_n)^{-1} = x_n^{-1} \cdots x_1^{-1}$, where $x_1, \dots, x_n \in X$. We denote by $\pi : X^* \rightarrow G$ the canonical projection, i.e., $\pi(u) = [u]_{\equiv}$. Let $\mathcal{A} = \langle \mathcal{X}, \mathcal{Q}, \delta, l, T \rangle$ be some finite automaton over X^* such that

$$\pi(L(\mathcal{A})) = K$$

is a subgroup of G . Without loss of generality, we can suppose that \mathcal{A} has a unique initial state q_0 and that \mathcal{A} is trim, i.e., that every state is accessible from $l = \{q_0\}$

and co-accessible from T . Let us define $\delta^{-1} = \{(r, x^{-1}, q) \mid (q, x, r) \in \delta\}$ and $\delta' = \delta \cup \delta^{-1}$. Since $\pi(\mathsf{L}(\mathcal{A})) = K$ is a subgroup of G , it follows easily that

$$\pi(\mathsf{L}(\mathcal{A})) = \pi(\mathsf{L}(\langle \mathcal{X}, \mathsf{Q}, \delta', \mathsf{l}, \mathsf{T} \rangle)).$$

Let us fix some subset $\lambda \subseteq \delta'$ which is a covering tree in the sense that for every $q \in \mathsf{Q}$, there exists a *unique* path $\Lambda(q)$ from q_0 to q , using only edges in λ . We denote by $W(q) \in X^*$ the sequence of labels of this path. Note that $W(q_0) = \varepsilon$. Let us define

$$\mathcal{G} = \{W(q)xW(r)^{-1} \mid (q, x, r) \in \delta\} \cup \{W(q) \mid q \in \mathsf{T}\} \subseteq X^*.$$

One can check that $K = \pi((\mathcal{G} \cup \mathcal{G}^{-1})^*)$. Thus, K is finitely generated. \square

The idea above is reminiscent of that of defining a set of generators for the fundamental group of a graph, see e.g. [Ser03, Section 5.1].

Definition 2 (Howson property). *A group G is said to have the Howson property if for every two finitely generated subgroups K_1 and K_2 of G , the subgroup $K_1 \cap K_2$ is finitely generated, too.*

Definition 3 (subgroup separation property, LERF). *A group G is said to have the subgroup separation property (also called local extended residual finiteness property — LERF for short) if for every finitely generated proper subgroup $K < G$ and every element $g \in G \setminus K$, there exists a subgroup Z which has finite index in G and such that $K \leq Z$ and $g \notin Z$.*

The LERF property has been investigated in several works, see e.g. [AG73, Git97, Hal49, MR04, Sco78, Wis00].

Remark 4. The following two conditions are easily seen to be equivalent to the LERF property for G , see e.g. [MR04, Wis00]:

- For every finitely generated proper subgroup $K < G$ and every element $g \in G \setminus K$, there exists a normal subgroup N which has finite index in G and such that $g \notin NK$.
- For every finitely generated proper subgroup $K < G$ and every element $g \in G \setminus K$, there exists a homomorphism $\pi : G \rightarrow S$ with S a finite group such that $\pi(g) \notin \pi(K)$.

2.3. Algorithmic problems

Let $\mathbb{M} = X^*/\equiv$ be a countably-generated monoid, where X is a *countable* generating set. Let $\pi : X^* \rightarrow X^*/\equiv$ be the canonical homomorphism. We assume \mathbb{M} to be countably generated in order to have a finite description of elements of \mathbb{M} by words over X .

The *word problem* for \mathbb{M} is the following decision problem:

instance: words $u, v \in X^*$

question: $\pi(u) = \pi(v)$?

Beside the word problem we shall investigate the following (more general) decision problems:

The *membership problem for rational subsets of \mathbb{M}* is the following decision problem:

instance: a finite automaton \mathcal{A} over X^* and a word $u \in X^*$.

question: $\pi(u) \in \pi(L(\mathcal{A}))$?

The *emptiness problem for boolean combinations of rational subsets of \mathbb{M}* is the following decision problem:

instance: a set $B \in \text{Bool}(\text{Rat}(\mathbb{M}))$ (represented by a boolean combination B of finite automata over X^*)

question: $B = \emptyset$?

Let us now assume that $\mathbb{M} = X^*/\equiv$ is a group. The *membership problem for finitely generated subgroups of \mathbb{M}* , also called the *generalized word problem* (GWP for short), is the following decision problem:

instance: words $u_1, \dots, u_n, u \in X^*$.

question: $\pi(u) \in \{\pi(u_1), \pi(u_1)^{-1}, \dots, \pi(u_n), \pi(u_n)^{-1}\}^*$, i.e., does $\pi(u)$ belong to the subgroup of \mathbb{M} generated by $\pi(u_1), \dots, \pi(u_n)$?

It is easy to see that the decidability status of all these decision problems is independent of the chosen generating set X for \mathbb{M} , i.e., they are indeed properties of \mathbb{M} .

3. HNN-extensions

Let us fix throughout this section a monoid H (the base monoid). In this paper, a subgroup of H is a submonoid of H , which forms a group.^a Equivalently, A is a subgroup of H if A is a subgroup of the group of units of H . Let us also fix two finite, isomorphic subgroups $A \leq H, B \leq H$ and an isomorphism $\varphi : A \rightarrow B$. We also fix some presentation $\langle X; \equiv_H \rangle$ of H , i.e., an alphabet X and a congruence \equiv_H over the free monoid X^* such that $H = X^*/\equiv_H$. Note that X is not supposed to be finite, in general. Without loss of generality we assume that every element $h \in A \cup B$ is of the form $[x]_{\equiv_H}$ for some letter $x \in X$. In particular, there is a letter 1 for the neutral element of H . Of course $1 \equiv_H \varepsilon$. We consider the HNN-extension

$$G = \langle H, t; t^{-1}at = \varphi(a)(a \in A) \rangle. \quad (7)$$

^aNote that usually, a subgroup A of H is defined to be a subsemigroup of H , which forms a group. In particular, the identity element of A may be an idempotent of H different from the identity of H . Nevertheless, we prefer to use the term subgroup for our definition in order to avoid too many different notions.

This means that $G = (X \cup \{t, t^{-1}\})^* / \equiv$ where \equiv is the smallest congruence over the free monoid $(X \cup \{t, t^{-1}\})^*$ such that:

$$\begin{aligned} \equiv_H &\subseteq \equiv \\ tt^{-1} &\equiv t^{-1}t \equiv \varepsilon \\ t^{-1}at &\equiv \varphi(a) \text{ for all } a \in A \end{aligned}$$

Let

$$\pi : (X \cup \{t, t^{-1}\})^* \rightarrow G \quad (8)$$

denote the canonical morphism defined by $\pi(w) = [w]_{\equiv}$ for $w \in (X \cup \{t, t^{-1}\})^*$. This morphism naturally factorizes as

$$\pi = \pi_t \circ \pi_G, \quad (9)$$

where

$$\pi_t : (X \cup \{t, t^{-1}\})^* \rightarrow H * \{t, t^{-1}\}^* \text{ and} \quad (10)$$

$$\pi_G : H * \{t, t^{-1}\}^* \rightarrow G. \quad (11)$$

The kernel of π_G coincides with the smallest congruence \approx over $H * \{t, t^{-1}\}^*$ such that:

$$tt^{-1} \approx t^{-1}t \approx 1 \quad (12)$$

$$at \approx t\varphi(a) \quad \text{for all } a \in A \quad (13)$$

$$bt^{-1} \approx t^{-1}\varphi^{-1}(b) \text{ for all } b \in B \quad (14)$$

Note that (12) and (13) together imply (14) but below we will need (13) and (14) without assuming (12). Clearly, $G = H * \{t, t^{-1}\}^* / \approx$.

An element of $s \in H * \{t, t^{-1}\}^*$ can be viewed as a word over the alphabet $H \cup \{t, t^{-1}\}$ which has the form

$$s = h_0 t^{\alpha_1} h_1 \cdots t^{\alpha_i} h_i \cdots t^{\alpha_n} h_n, \quad (15)$$

where $n \geq 0$, $\alpha_i \in \{1, -1\}$, and $h_i \in H$. Such an element $s \in H * \{t, t^{-1}\}^*$ is also called a *t-sequence*. The *t-sequence* s is said to be a *reduced sequence* if it neither contains a factor of the form $t^{-1}at$ (with $a \in A$) nor tbt^{-1} (with $b \in B$). Let

$$\text{Red}(H, t) = \{s \in H * \{t, t^{-1}\}^* \mid s \text{ is reduced}\}.$$

We define the binary relation $\overset{1}{\sim}$ over $H * \{t, t^{-1}\}^*$ as follows: For all $s, s' \in H * \{t, t^{-1}\}^*$ let

$$\begin{aligned} s \overset{1}{\sim} s' &\Leftrightarrow \exists s_1, s_2 \in H * \{t, t^{-1}\}^*, a \in A, b \in B : \\ &\left\{ \begin{array}{l} (s = s_1 t s_2 \wedge s' = s_1 a^{-1} t \varphi(a) s_2) \vee \\ (s = s_1 t^{-1} s_2 \wedge s' = s_1 b^{-1} t^{-1} \varphi^{-1}(b) s_2) \end{array} \right\}. \end{aligned}$$


Let us denote by \sim the congruence over $H * \{t, t^{-1}\}^*$ generated by all the rules of type (13) and (14) above. It coincides with the reflexive and transitive closure of $\overset{1}{\sim}$. Equivalently, if

$$\begin{aligned} s &= h_0 t^{\alpha_1} h_1 \cdots t^{\alpha_n} h_n \text{ and} \\ s' &= h'_0 t^{\alpha'_1} h'_1 \cdots t^{\alpha'_m} h'_m \end{aligned}$$

(with $n, m \geq 0$, $\alpha_i, \alpha'_j \in \{1, -1\}$ and $h_i, h'_j \in H$), then $s \sim s'$ if and only if $n = m$, $\alpha_i = \alpha'_i$ for $1 \leq i \leq n$, and there exist $c_1, \dots, c_{2n} \in A \cup B$ such that:

- if $\alpha_i = 1$ then $c_{2i-1} \in A$ and $c_{2i} = \varphi(c_{2i-1}) \in B$ ($1 \leq i \leq n$)
- if $\alpha_i = -1$ then $c_{2i} \in A$ and $c_{2i-1} = \varphi(c_{2i}) \in B$ ($1 \leq i \leq n$)
- $h_i c_{2i+1} = c_{2i} h'_i$ in H for $0 \leq i \leq n$ (here we set $c_0 = c_{2n+1} = 1$)

This situation can be visualized by a diagram of the following form (also called a Van Kampen diagram, see [LS77] for more details), where $n = m = 4$. Light-shaded (resp. dark-shaded) areas represent relations in H (resp. relations of the form $at = t\varphi(a)$ ($a \in A$) or $bt^{-1} = t^{-1}\varphi^{-1}(b)$ ($b \in B$)).



The elements c_1, \dots, c_{2n} in such a diagram are also called *connecting elements*.

The set $\text{Red}(H, t)$ is saturated by the congruence \sim , i.e., $s \sim s'$ implies $s \in \text{Red}(H, t) \Leftrightarrow s' \in \text{Red}(H, t)$: just notice that, since A and B are groups, $aha' \in A \Leftrightarrow h \in A$ for all $h \in H, a, a' \in A$ and $bhb' \in B \Leftrightarrow h \in B$ for all $h \in H, b, b' \in B$. This property would fail if A and B were assumed to be merely submonoids of H .

One has $\sim \subseteq \approx$. Moreover, for reduced sequences the following fundamental lemma holds.

Lemma 5. *Let $s, s' \in \text{Red}(H, t)$. Then $s \approx s'$ if and only if $s \sim s'$.*

In the case where H is a group, Lemma 5 is an immediate consequence of [LS77, Theorem 2.1, p.182]. The extension to monoids can be proved in the same way as for groups using an idea of Artin and Van der Waerden (see Appendix A for a detailed verification). Lemma 5 implies that H is a submonoid of G . Another consequence is:

Corollary 6. *If H has a decidable word problem, then also has G a decidable word problem.*

Proof. Let $w, w' \in (X \cup \{t, t^{-1}\})^*$ (recall that X is a monoid generating set for H). We describe a procedure for checking whether $w \equiv w'$ or not. The words w and

w' have unique decompositions

$$w = u_0 t^{\alpha_1} u_1 \cdots t^{\alpha_i} u_i \cdots t^{\alpha_n} u_n \text{ and } w' = u'_0 t^{\alpha'_1} u'_1 \cdots t^{\alpha'_j} u'_j \cdots t^{\alpha'_m} u'_m,$$

where $n, m \geq 0, u_i, u'_j \in X^*$, and $\alpha_i, \alpha'_j \in \{1, -1\}$. Let the semi-Thue system ST over the alphabet $X \cup \{t, t^{-1}\}$ consist of the following rules:

$$t^{-1}ut \rightarrow \varphi([u]_{\equiv_H}) \quad \text{for } [u]_{\equiv_H} \in A \quad (16)$$

$$tut^{-1} \rightarrow \varphi^{-1}([u]_{\equiv_H}) \quad \text{for } [u]_{\equiv_H} \in B \quad (17)$$

Note that $\text{Red}(H, t) = \pi_t(\text{Irr}(\text{ST}))$. Since the word problem for H is decidable and A and B are finite, we can effectively compute from w and w' words z and z' such that $w \rightarrow_{\text{ST}}^* z \in \text{Irr}(\text{ST})$ and $w' \rightarrow_{\text{ST}}^* z' \in \text{Irr}(\text{ST})$. Assume that

$$z = v_0 t^{\beta_1} v_1 \cdots t^{\beta_i} v_i \cdots t^{\beta_p} v_p \text{ and } z' = v'_0 t^{\beta'_1} v'_1 \cdots t^{\beta'_i} v'_i \cdots t^{\beta'_q} v'_q, \quad (18)$$

where $p, q \geq 0, v_i, v'_j \in X^*$, and $\beta_i, \beta'_j \in \{1, -1\}$. Note that $\rightarrow_{\text{ST}} \subseteq \equiv$. Hence,

$$\begin{aligned} w \equiv w' &\Leftrightarrow z \equiv z' \\ &\Leftrightarrow \pi_G(\pi_t(z)) = \pi_G(\pi_t(z')) \\ &\Leftrightarrow \pi_t(z) \approx \pi_t(z') \\ &\Leftrightarrow \pi_t(z) \sim \pi_t(z'), \end{aligned}$$

where the last equivalence holds due to Lemma 5 and the fact that $\pi_t(z), \pi_t(z') \in \pi_t(\text{Irr}(\text{ST})) = \text{Red}(H, t)$. If $p \neq q$ or ($p = q$ and $\beta_i \neq \beta'_i$ for some $1 \leq i \leq p$) in (18), then $\pi_t(z) \not\sim \pi_t(z')$ and thus $w \not\equiv w'$. Hence, assume that

$$z = v_0 t^{\beta_1} v_1 \cdots t^{\beta_i} v_i \cdots t^{\beta_p} v_p \text{ and } z' = v'_0 t^{\beta_1} v'_1 \cdots t^{\beta_i} v'_i \cdots t^{\beta_p} v'_p.$$

We now guess the connecting elements between $\pi_t(z)$ and $\pi_t(z')$ in a diagram of the form (\dagger) . More precisely, we nondeterministically guess elements $c_1, \dots, c_{2n} \in A \cup B$ such that:

- if $\beta_i = 1$ then $c_{2i-1} \in A$ and $c_{2i} = \varphi(c_{2i-1}) \in B$ ($1 \leq i \leq p$) and
- if $\beta_i = -1$ then $c_{2i} \in A$ and $c_{2i-1} = \varphi(c_{2i}) \in B$ ($1 \leq i \leq p$).

Since A and B are finite, this guessing can be done by a nondeterministic Turing machine. Finally, using the decidability of the word problem for H , we check whether $v_i c_{2i+1} \equiv_H c_{2i} v'_i$ for $0 \leq i \leq p$ (here we set $c_0 = c_{2p+1} = 1$). If this is true for some choice of $c_1, \dots, c_{2n} \in A \cup B$, then we know that indeed $w \equiv w'$. \square

3.1. HNN-extensions and morphisms

Suppose we are given a monoid homomorphism: $\pi' : H \rightarrow H'$ fulfilling

$$\text{Ker}(\pi') \cap A \times A = \text{Id}_A \quad \text{and} \quad \text{Ker}(\pi') \cap B \times B = \text{Id}_B. \quad (19)$$

Thus, A and B are subgroups of H' . Hence, we can define the HNN-extension

$$G' = \langle H', t; t^{-1}at = \varphi(a)(a \in A) \rangle. \quad (20)$$

$$\begin{array}{ccc}
 H * \{t, t^{-1}\}^* & \xrightarrow{\pi'_*} & H' * \{t, t^{-1}\}^* \\
 \downarrow \pi_G & & \downarrow \pi_{G'} \\
 G & \xrightarrow{\pi'} & G'
 \end{array}$$

 Fig. 1. Maps induced by π'

In the presentation (20), every relation $t^{-1}at = \varphi(a)$ should be written as $t^{-1}\pi'(a)t = \pi'(\varphi(a))$, but (19) allows us to identify every $a \in A$ with its image $\pi'(a) \in \pi'(A)$. The map π' induces a map $\pi'_* : H * \{t, t^{-1}\}^* \rightarrow H' * \{t, t^{-1}\}^*$, which is the unique monoid homomorphism such that

$$\forall h \in H : \pi'_*(h) = \pi'(h), \quad \pi'_*(t) = t, \quad \pi'_*(t^{-1}) = t^{-1}.$$

The composite morphism $\pi'_* \circ \pi_{G'} : H * \{t, t^{-1}\}^* \rightarrow G'$ has a kernel which contains the congruence \approx . Hence, it induces a map $\tilde{\pi}' : G \rightarrow G'$ such that

$$\pi'_* \circ \pi_{G'} = \pi_G \circ \tilde{\pi}'.$$

One can check that

$$\forall h \in H : \tilde{\pi}'(h) = \pi'(h), \quad \tilde{\pi}'(t) = t, \quad \tilde{\pi}'(t^{-1}) = t^{-1}.$$

Hence $\tilde{\pi}'$ is an extension of π' , therefore we shall (abusively) also use the symbol π' to denote $\tilde{\pi}'$. We represent the situation by the commutative diagram in Figure 1.

4. Finite automata for HNN-extensions

Let us fix throughout this section an HNN-extension G of a base monoid H as described by (7) with A and $B = \varphi(A)$ finite. We define below a special kind of finite automata, which will be well-suited for our aim of showing that some decidability and structural properties of rational subsets of H are transferred to G .

4.1. Finite t -automata

We define here a kind of finite automata that recognize subsets of $H * \{t, t^{-1}\}^*$ via transitions that either read one of the letters t, t^{-1} or read an element of H , provided it belongs to some definite set $F \in \mathcal{F}$. The typical class \mathcal{F} that we have in mind is the set $\text{Rat}(H)$ of all rational subsets of H , or the set \mathcal{S} of all right-cosets of finitely generated subgroups of H (in case H is a group), or some other families of subsets of H derived in a natural way from the two above families.

Let $\mathcal{F} \subseteq \mathcal{P}(H)$ be a set of subsets of H such that

$$\forall c \in A \cup B : \{c\} \in \mathcal{F}.$$

A *finite t -automaton*, briefly *fta*, over $H * \{t, t^{-1}\}^*$ with labelling set \mathcal{F} is a 5-tuple

$$\mathcal{A} = \langle \mathcal{L}, \mathcal{Q}, \delta, \mathbf{l}, \mathbf{T} \rangle, \tag{21}$$

where

$$\mathcal{L} \subseteq \mathcal{F} \cup \{\{t\}, \{t^{-1}\}\}$$

is finite, \mathbf{Q} is a finite set of states, $\mathbf{I} \subseteq \mathbf{Q}$ is the set of initial states, $\mathbf{T} \subseteq \mathbf{Q}$ is the set of terminal states, and δ (the set of transitions) is a subset of $\mathbf{Q} \times \mathcal{L} \times \mathbf{Q}$ such that

$$\forall q \in \mathbf{Q} \exists L \subseteq \mathcal{L} : 1 \in L \wedge (q, L, q) \in \delta. \quad (23)$$

We define

$$\widehat{\delta} = \{(p, h, q) \in \mathbf{Q} \times (H \cup \{t, t^{-1}\}) \times \mathbf{Q} \mid \exists L \subseteq \mathcal{L} : h \in L \wedge (p, L, q) \in \delta\}. \quad (24)$$

The automaton \mathcal{A} induces a representation map

$$\mu_{\mathcal{A}} : H * \{t, t^{-1}\}^* \rightarrow \text{BR}(\mathbf{Q})$$

defined as follows: First, define $\mu_{\mathcal{A},0} : H \cup \{t, t^{-1}\} \rightarrow \text{BR}(\mathbf{Q})$ as follows, where $h \in H \cup \{t, t^{-1}\}$:

$$\mu_{\mathcal{A},0}(h) = \{(q, r) \in \mathbf{Q} \times \mathbf{Q} \mid (q, h, r) \in \widehat{\delta}\} \quad (25)$$

(note that $\text{Id}_{\mathbf{Q}} \subseteq \mu_{\mathcal{A},0}(1)$ due to (23)). For $h \in H \cup \{t, t^{-1}\}$ and $q, r \in \mathbf{Q}$ we will also write $q \xrightarrow{h}_{\mathcal{A}} r$ instead of $(q, r) \in \mu_{\mathcal{A},0}(h)$.

Now, let $s \in H * \{t, t^{-1}\}^*$ be a sequence of the form (15). Then

$$\begin{aligned} \mu_{\mathcal{A}}(s) &= \mu_{\mathcal{A},0}(h_0) \circ \mu_{\mathcal{A},0}(t^{\alpha_1}) \circ \mu_{\mathcal{A},0}(h_1) \cdots \\ &\quad \mu_{\mathcal{A},0}(t^{\alpha_i}) \circ \mu_{\mathcal{A},0}(h_i) \cdots \mu_{\mathcal{A},0}(t^{\alpha_n}) \circ \mu_{\mathcal{A},0}(h_n). \end{aligned} \quad (26)$$

Notice that for the definition of the map $\mu_{\mathcal{A}}$ we only consider paths in \mathcal{A} , which alternate strictly between H -edges and $t^{\pm 1}$ -edges. As a consequence of this definition, the second property stated in (6) need not be fulfilled by this kind of automata in case $m, m' \in H$. Instead of $(q, r) \in \mu_{\mathcal{A}}(s)$ we also write $q \xrightarrow{s}_{\mathcal{A}} r$ (since $\mu_{\mathcal{A},0}(h) = \mu_{\mathcal{A}}(h)$ for $h \in H \cup \{t, t^{-1}\}$, this notation extends our notation introduced earlier).

The subset of $H * \{t, t^{-1}\}^*$ recognized by \mathcal{A} is



$$\text{L}(\mathcal{A}) = \{s \in H * \{t, t^{-1}\}^* \mid \mu_{\mathcal{A}}(s) \cap (\mathbf{I} \times \mathbf{T}) \neq \emptyset\}.$$

Let us introduce a finite set \mathcal{T}_6 of 6 types, which will induce a partition of the states of an fta:

$$\mathcal{T}_6 = \{(A, T), (B, T), (1, H), (1, 1), (A, H), (B, H)\} \quad (27)$$

The first component of the types will serve below for defining a normal form for automata; the second component suggests the possible labels going out of a state of that type. We define a directed, edge-labeled graph $\mathcal{G}_6 = (\mathcal{T}_6, \mathcal{E}_6)$ by

$$\begin{aligned} \mathcal{E}_6 &= \{((C, D), C, (C, D)) \mid (C, D) \in \mathcal{T}_6\} \\ &\cup \{((A, T), t, (B, H)), ((B, T), t^{-1}, (A, H))\} \\ &\cup \{((C, H), H, (C, T)), ((1, H), H, (C, T)) \mid C \in \{A, B\}\} \\ &\cup \{((C, H), H, (1, 1)) \mid C \in \{1, A, B\}\} \\ &\cup \{((B, H), H, (A, T)), ((A, H), H, (B, T))\}. \end{aligned}$$


 Fig. 2. The graph $\mathcal{G}_6 = (\mathcal{T}_6, \mathcal{E}_6)$

 Fig. 3. The graph $\mathcal{R}_6 = (\mathcal{T}_6, \mathcal{E}'_6)$

The graph \mathcal{G}_6 is represented in Figure 2. We sometimes use also the graph $\mathcal{R}_6 = (\mathcal{T}_6, \mathcal{E}'_6)$ where

$$\mathcal{E}'_6 = \mathcal{E}_6 \setminus \{((A, H), H, (A, T)), ((B, H), H, (B, T))\} \cup \{((A, H), H \setminus A, (A, T)), ((B, H), H \setminus B, (B, T))\};$$

it is shown in Figure 3. One can check that the graph \mathcal{G}_6 (resp. \mathcal{R}_6) endowed with the set of initial states $\mathsf{l}_6 = \{(1, H)\}$ and the set of final states $\mathsf{T}_6 = \{(1, 1)\}$ is an fta recognizing $H * \{t, t^{-1}\}^*$ (resp. $\text{Red}(H, t)$).^b For this, the A -loops and B -loops are not really relevant, but these loops will be useful later, when we introduce the property of \sim -saturation, see Definition 9.

Since \mathcal{G}_6 and \mathcal{R}_6 contain only a single t -transition (resp. t^{-1} -transition), one can easily check the following (recall the definition of the $\hat{\cdot}$ -operator from (24)):

Lemma 7. *For every sequence $s \in H * \{t, t^{-1}\}^*$ of the form (15) there is exactly one sequence $\Theta_0, \Theta_1, \dots, \Theta_{2n+1} \in \mathcal{T}_6$ of types such that*

^bFormally, the label t^α ($\alpha \in \{1, -1\}$) of \mathcal{G}_6 (resp. \mathcal{R}_6) has to be replaced by a transition labelled with the singleton set $\{t^\alpha\}$ in order to make \mathcal{G}_6 (resp. \mathcal{R}_6) an fta. In the rest of this paper, we will repeatedly write t^α , when we formally have to write $\{t^\alpha\}$.

- (1) $\Theta_0 = (1, H)$, $\Theta_{2n+1} = (1, 1)$,
- (2) $(\Theta_{2i}, h_i, \Theta_{2i+1}) \in \widehat{\mathcal{E}}_6$ for all $0 \leq i \leq n$, and
- (3) $(\Theta_{2i-1}, t_i^{\alpha_i}, \Theta_{2i}) \in \widehat{\mathcal{E}}_6$ for all $1 \leq i \leq n$.

Moreover, we must have $\Theta_i \neq \Theta_{i+1}$ for all $0 \leq i \leq 2n$.

Lemma 7 implies that that every sequence s can be accepted in a unique way by the fta \mathcal{G}_6 . The same statement also holds for \mathcal{R}_6 if we restrict to reduced sequences.

Definition 8 (partitioned fta). A partitioned fta with labelling set \mathcal{F} is a 6-tuple

$$\mathcal{A} = \langle \mathcal{L}, \mathcal{Q}, \tau, \delta, \mathsf{l}, \mathsf{T} \rangle, \quad (28)$$

where \mathcal{L} , \mathcal{Q} , δ , l , and T are as in (21), and $\tau : \mathcal{Q} \rightarrow \mathcal{T}_6$ assigns a type to every state, and the transitions of δ and the sets l and T respect the types in the following sense:

$$\forall (q, h, r) \in \widehat{\delta} : (\tau(q), h, \tau(r)) \in \widehat{\mathcal{E}}_6 \quad (29)$$

$$\tau(\mathsf{l}) = \{(1, H)\} \wedge \tau(\mathsf{T}) = \{(1, 1)\}. \quad (30)$$

Recall the equivalences \sim and \approx over $H * \{t, t^{-1}\}^*$ defined in Section 3.

Definition 9 (\approx -compatible, \sim -saturated). An fta \mathcal{A} is said to be \approx -compatible if and only if

$$[\mathsf{L}(\mathcal{A})]_{\approx} = [\mathsf{L}(\mathcal{A}) \cap \text{Red}(H, t)]_{\approx}. \quad (31)$$

It is said to be \sim -saturated if and only if

$$\forall s, s' \in H * \{t, t^{-1}\}^* : s \sim s' \Rightarrow \mu_{\mathcal{A}}(s) = \mu_{\mathcal{A}}(s'). \quad (32)$$

Remark 10. The following observations are easy to verify:

- Property (31) is equivalent to $\mathsf{L}(\mathcal{A}) \subseteq [\mathsf{L}(\mathcal{A}) \cap \text{Red}(H, t)]_{\approx}$, i.e., for every sequence $s \in \mathsf{L}(\mathcal{A})$ there must exist a reduced sequence $s' \in \mathsf{L}(\mathcal{A})$ such that $s \approx s'$.
- If $\mathsf{L}(\mathcal{A}) \subseteq \text{Red}(H, t)$, then \mathcal{A} is \approx -compatible.
- \mathcal{A} is \sim -saturated if and only if for all states $q_1, q_2, q_3, q_4 \in \mathcal{Q}$, all $h_1, h_2 \in H$, and all $a \in A$, $b \in B$ such that

$$\begin{aligned} & q_1 \xrightarrow{h_1}_{\mathcal{A}} q_2 \xrightarrow{t}_{\mathcal{A}} q_3 \xrightarrow{h_2}_{\mathcal{A}} q_4 \\ \text{(resp. } & q_1 \xrightarrow{h_1}_{\mathcal{A}} q_2 \xrightarrow{t^{-1}}_{\mathcal{A}} q_3 \xrightarrow{h_2}_{\mathcal{A}} q_4), \end{aligned}$$

there exist $q'_2, q'_3 \in \mathcal{Q}$ with

$$\begin{aligned} & q_1 \xrightarrow{h_1 a^{-1}}_{\mathcal{A}} q'_2 \xrightarrow{t}_{\mathcal{A}} q'_3 \xrightarrow{\varphi(a)h_2}_{\mathcal{A}} q_4 \\ \text{(resp. } & q_1 \xrightarrow{h_1 b^{-1}}_{\mathcal{A}} q'_2 \xrightarrow{t^{-1}}_{\mathcal{A}} q'_3 \xrightarrow{\varphi^{-1}(b)h_2}_{\mathcal{A}} q_4). \end{aligned}$$

- Whereas \approx -compatibility depends on $L(\mathcal{A})$ only, \sim -saturation does really depend on the automaton \mathcal{A} itself.
- If \mathcal{A} is \sim -saturated, then the set $L(\mathcal{A})$ is \sim -saturated, i.e., if $s \sim s'$, then $s \in L(\mathcal{A})$ if and only if $s' \in L(\mathcal{A})$.
- The partitioned fta \mathcal{G}_6 and \mathcal{R}_6 are both \approx -compatible and \sim -saturated. Since we require (23), the A -labeled (resp. B -labeled) loops at state (A, T) and (A, H) (resp. (B, T) and (B, H)) in Figure 2 and 3 are crucial for \sim -saturations of \mathcal{G}_6 and \mathcal{R}_6 .

Definition 11 (unitary fta). *The partitioned fta $\mathcal{A} = \langle \mathcal{L}, \mathcal{Q}, \tau, \delta, \mathfrak{l}, \mathfrak{T} \rangle$ is said to be unitary if and only if for every type $\Theta \in \mathcal{T}_6$ we have*

$$\mu_{\mathcal{A},0}(1) \cap (\tau^{-1}(\Theta) \times \tau^{-1}(\Theta)) = \text{Id}_{\mathcal{Q}} \cap (\tau^{-1}(\Theta) \times \tau^{-1}(\Theta)).$$

This last technical condition turns out to be useful in [LS06a]. It means that no edge between two different states of the same type is labelled with a set containing 1.

Definition 12 (deterministic fta). *The partitioned fta $\mathcal{A} = \langle \mathcal{L}, \mathcal{Q}, \tau, \delta, \mathfrak{l}, \mathfrak{T} \rangle$ is said to be deterministic if and only if*

$$\text{Card}(\mathfrak{l}) = 1 \tag{33}$$

(i.e. it has exactly one initial state) and for all $q, r, r' \in \mathcal{Q}$ and all $L, L' \in \mathcal{L}$,

$$((q, L, r), (q, L', r') \in \delta \wedge L \cap L' \neq \emptyset \wedge \tau(r) = \tau(r')) \Rightarrow (L = L' \wedge r = r'). \tag{34}$$

Clearly, \mathcal{G}_6 and \mathcal{R}_6 are both deterministic, since these fta both have only one initial state and for every type there exists only one state of that type.

Finally, we define the notion of a complete partitioned fta. The intuitive idea is that of a local condition over the labels of transitions, which ensures that for every reduced sequence $s \in \text{Red}(H, t)$ and every state q of type $(1, H)$, we have $q \xrightarrow{s}_{\mathcal{A}} r$ for some state r of type $(1, 1)$. For every $q \in \mathcal{Q}$ and every $\Theta \in \mathcal{T}_6$ we thus introduce the set

$$L^1(q, \Theta) = \bigcup \{L \mid \exists r \in \tau^{-1}(\Theta) : (q, L, r) \in \delta\}. \tag{35}$$

Definition 13 (complete fta). *The partitioned fta $\mathcal{A} = \langle \mathcal{L}, \mathcal{Q}, \tau, \delta, \mathfrak{l}, \mathfrak{T} \rangle$ is said to be complete if and only if*

$$\forall (\Theta, C, \Theta') \in \mathcal{E}'_6 \forall q \in \tau^{-1}(\Theta) : C \subseteq L^1(q, \Theta').$$

Note that \mathcal{G}_6 and \mathcal{R}_6 are both complete.

4.2. Operations on fta

In this section we introduce operations on fta, which are analogues of the classical complement and product construction for finite automata.

Definition 14 (product automaton). *Given two fta $\mathcal{A}_i = \langle \mathcal{L}_i, \mathbf{Q}_i, \delta_i, \mathbf{l}_i, \mathbf{T}_i \rangle$ ($i \in \{1, 2\}$), we define the product fta $\mathcal{A}_1 \times \mathcal{A}_2 = \langle \mathcal{L}, \mathbf{Q}, \delta, \mathbf{l}, \mathbf{T} \rangle$, where:*

$$\begin{aligned} \mathcal{L} &= \{L_1 \cap L_2 \mid L_1 \in \mathcal{L}_1, L_2 \in \mathcal{L}_2\} \setminus \{\emptyset\} \\ \mathbf{Q} &= \mathbf{Q}_1 \times \mathbf{Q}_2, \\ \mathbf{l} &= \mathbf{l}_1 \times \mathbf{l}_2 \\ \mathbf{T} &= \mathbf{T}_1 \times \mathbf{T}_2 \\ \delta &= \{(p_1, p_2), L_1 \cap L_2, (q_1, q_2) \mid (p_1, L_1, q_1) \in \delta_1, (p_2, L_2, q_2) \in \delta_2\} \end{aligned}$$

For two partitioned fta $\mathcal{A}_i = \langle \mathcal{L}_i, \mathbf{Q}_i, \tau_i, \delta_i, \mathbf{l}_i, \mathbf{T}_i \rangle$ the product automaton is the partitioned fta $\mathcal{A}_1 \times \mathcal{A}_2 = \langle \mathcal{L}, \mathbf{Q}, \tau, \delta, \mathbf{l}, \mathbf{T} \rangle$, where \mathcal{L}, \mathbf{l} , and \mathbf{T} are defined as above, but

$$\mathbf{Q} = \bigcup_{\Theta \in \mathcal{T}_6} \tau_1^{-1}(\Theta) \times \tau_2^{-1}(\Theta),$$

$$\delta = \{(p_1, p_2), L_1 \cap L_2, (q_1, q_2) \mid (p_1, L_1, q_1) \in \delta_1, (p_2, L_2, q_2) \in \delta_2\} \cap \mathbf{Q} \times \mathcal{L} \times \mathbf{Q},$$

and $\tau(p_1, p_2) = \tau_1(p_1) = \tau_2(p_2)$ for $(p_1, p_2) \in \mathbf{Q}$.

Lemma 15. *For two fta (resp. partitioned fta) \mathcal{A}_1 and \mathcal{A}_2 we have $L(\mathcal{A}_1 \times \mathcal{A}_2) = L(\mathcal{A}_1) \cap L(\mathcal{A}_2)$.*

Proof. For ordinary (non-partitioned) fta the statement of the lemma is obvious. For partitioned fta one has to use Lemma 7. It implies that if $s \in L(\mathcal{A}_1) \cap L(\mathcal{A}_2)$, then the state sequences in accepting runs of \mathcal{A}_1 and \mathcal{A}_2 , respectively, on s are mapped to the same sequence of types via the type functions of \mathcal{A}_1 and \mathcal{A}_2 , respectively. \square

Lemma 15 implies:

Lemma 16. *Let \mathcal{A} be some fta with labelling set \mathcal{F} . The fta $\mathcal{A} \times \mathcal{G}_6$ endowed with the map $\tau : (p, \Theta) \mapsto \Theta$ is a partitioned fta with labelling set \mathcal{F} and $L(\mathcal{A} \times \mathcal{G}_6) = L(\mathcal{A})$.*

Lemma 17. *Let \mathcal{A}_1 and \mathcal{A}_2 be fta (resp. partitioned fta). Then we have:*

- (a) *If \mathcal{A}_1 and \mathcal{A}_2 are both \sim -saturated, then $\mathcal{A}_1 \times \mathcal{A}_2$ is \sim -saturated, too.*
- (b) *If \mathcal{A}_1 and \mathcal{A}_2 are both \approx -compatible and \sim -saturated, then $\mathcal{A}_1 \times \mathcal{A}_2$ is \approx -compatible and \sim -saturated, too.*

Proof. For statement (a) and non-partitioned fta assume that in the product automaton $\mathcal{A} = \mathcal{A}_1 \times \mathcal{A}_2$ we have

$$(q_1, p_1) \xrightarrow{h_1}_{\mathcal{A}} (q_2, p_2) \xrightarrow{t}_{\mathcal{A}} (q_3, p_3) \xrightarrow{h_2}_{\mathcal{A}} (q_4, p_4).$$

Hence,

$$\begin{aligned} q_1 &\xrightarrow{\mathcal{A}_1} h_1 q_2 \xrightarrow{\mathcal{A}_1} t q_3 \xrightarrow{\mathcal{A}_1} h_2 q_4 \text{ and} \\ p_1 &\xrightarrow{\mathcal{A}_2} h_1 p_2 \xrightarrow{\mathcal{A}_2} t p_3 \xrightarrow{\mathcal{A}_2} h_2 p_4. \end{aligned}$$

Since \mathcal{A}_1 and \mathcal{A}_2 are both \sim -saturated, we have, for every $a \in A$,

$$\begin{aligned} q_1 &\xrightarrow{\mathcal{A}_1} h_1 a^{-1} q'_2 \xrightarrow{\mathcal{A}_1} t q'_3 \xrightarrow{\mathcal{A}_1} \varphi(a) h_2 q_4 \text{ and} \\ p_1 &\xrightarrow{\mathcal{A}_2} h_1 a^{-1} p'_2 \xrightarrow{\mathcal{A}_2} t p'_3 \xrightarrow{\mathcal{A}_2} \varphi(a) h_2 p_4 \end{aligned}$$

for states q'_2 and q'_3 (resp. p'_2 and p'_3) of \mathcal{A}_1 (resp. \mathcal{A}_2). Hence,

$$(q_1, p_1) \xrightarrow{\mathcal{A}} h_1 a^{-1} (q'_2, p'_2) \xrightarrow{\mathcal{A}} t (q'_3, p'_3) \xrightarrow{\mathcal{A}} \varphi(a) h_2 (q_4, p_4).$$

An analogous argument for t^{-1} shows that $\mathcal{A} = \mathcal{A}_1 \times \mathcal{A}_2$ is indeed \sim -saturated. For partitioned fta we can reason similarly, we just have to note that the states q_2, q'_2, p_2, p'_2 (resp. q_3, q'_3, p_3, p'_3) have to be of type (A, T) (resp. (B, H)). This, ensures that (q'_2, p'_2) and (q'_3, p'_3) are states of the partitioned fta \mathcal{A} .

For (b) it suffices by Lemma 15 and statement (a) to prove that $L(\mathcal{A}_1) \cap L(\mathcal{A}_2) \subseteq [L(\mathcal{A}_1) \cap L(\mathcal{A}_2) \cap \text{Red}(H, t)]_{\approx}$. So, assume that

$$s \in L(\mathcal{A}_1) \cap L(\mathcal{A}_2).$$

Since \mathcal{A}_1 and \mathcal{A}_2 are both \approx -compatible, we have

$$s \in [L(\mathcal{A}_1) \cap \text{Red}(H, t)]_{\approx} \cap [L(\mathcal{A}_2) \cap \text{Red}(H, t)]_{\approx}.$$

Thus, there exist sequences s_1 and s_2 such that

$$s \approx s_1 \in L(\mathcal{A}_1) \cap \text{Red}(H, t) \text{ and } s \approx s_2 \in L(\mathcal{A}_2) \cap \text{Red}(H, t).$$

Hence, $s_1 \approx s_2$ and Lemma 5 implies $s_1 \sim s_2 \in L(\mathcal{A}_2)$. Since \mathcal{A}_2 is \sim -saturated, we have $s_1 \in L(\mathcal{A}_2)$. Thus,

$$s \approx s_1 \in L(\mathcal{A}_1) \cap L(\mathcal{A}_2) \cap \text{Red}(H, t),$$

i.e., $s \in [L(\mathcal{A}_1) \cap L(\mathcal{A}_2) \cap \text{Red}(H, t)]_{\approx}$. □

The next lemma is straightforward to prove.

Lemma 18. *Let \mathcal{A}_1 and \mathcal{A}_2 be partitioned fta. Then we have:*

- (a) *If \mathcal{A}_1 and \mathcal{A}_2 are both deterministic, then $\mathcal{A}_1 \times \mathcal{A}_2$ is deterministic, too.*
- (b) *If \mathcal{A}_1 and \mathcal{A}_2 are both complete, then $\mathcal{A}_1 \times \mathcal{A}_2$ is complete, too.*

Definition 19 (complement automaton). *For a partitioned, deterministic, and complete fta $\mathcal{A} = \langle \mathcal{L}, \mathcal{Q}, \tau, \delta, l, \top \rangle$ we define the complement fta $\bar{\mathcal{A}} = \langle \mathcal{L}, \mathcal{Q}, \tau, \delta, l, \tau^{-1}(1, 1) \setminus \top \rangle \times \mathcal{R}_6$. It is again a partitioned, deterministic, and complete fta (by Lemma 18 and the fact that \mathcal{R}_6 is deterministic and complete).*

Lemma 20. *Let \mathcal{A} be a partitioned, deterministic, and complete fta. Then $L(\overline{\mathcal{A}}) = \text{Red}(H, t) \setminus L(\mathcal{A})$. Moreover, if \mathcal{A} is \sim -saturated, then $\overline{\mathcal{A}}$ is again \sim -saturated.*

Proof. For the first statement, note that $L(\overline{\mathcal{A}}) \subseteq \text{Red}(H, t)$ since we take the product with \mathcal{R}_6 . Moreover, for a partitioned, deterministic, and complete fta \mathcal{A} , there exists for every reduced sequence $s \in \text{Red}(H, t)$ exactly one state $q \in \tau_{\mathcal{A}}^{-1}(1, 1)$ such that $(q_0, q) \in \mu_{\mathcal{A}}(s)$, where q_0 is the unique initial state of \mathcal{A} . This implies $L(\overline{\mathcal{A}}) = \text{Red}(H, t) \setminus L(\mathcal{A})$.

For the second statement, note that $\langle \mathcal{L}, \mathcal{Q}, \tau, \delta, l, \tau^{-1}(1, 1) \setminus \mathbb{T} \rangle$ is \sim -saturated, since this automaton has the same μ -mapping as the \sim -saturated fta \mathcal{A} . Hence, $\overline{\mathcal{A}}$ is \sim -saturated by Lemma 17 and the fact that \mathcal{R}_6 is \sim -saturated. \square

4.3. Automata and morphisms

Let us consider another HNN-extension G' defined by an homomorphism $\pi' : H \rightarrow H'$ satisfying (19) as described in Section 3.1 by (20). Given an fta $\mathcal{A} = \langle \mathcal{L}, \mathcal{Q}, \delta, l, \mathbb{T} \rangle$, we define

$$\pi'_*(\mathcal{A}) = \langle \mathcal{L}', \mathcal{Q}, \delta', l, \mathbb{T} \rangle \quad (36)$$

with $\mathcal{L}' = \{\pi'(L) \mid L \in \mathcal{L}\}$ and

$$\delta' = \{(q, \pi'(L), r) \in \mathcal{Q} \times \mathcal{L}' \times \mathcal{Q} \mid (q, L, r) \in \delta\}.$$

We denote the congruence on $H' * \{t, t^{-1}\}$, which is defined in the same way as \sim on $H * \{t, t^{-1}\}$, with \sim' .

Lemma 21. *Let \mathcal{A} be some fta. Then we have:*

- (a) $L(\pi'_*(\mathcal{A})) = \pi'_*(L(\mathcal{A}))$
- (b) *If \mathcal{A} is \sim -saturated, then $\pi'_*(\mathcal{A})$ is \sim' -saturated.*

Proof. Statement (a) is obvious. For statement (b) assume that \mathcal{A} is \sim -saturated and that

$$q_1 \xrightarrow{h_1} \pi'_*(\mathcal{A}) q_2 \xrightarrow{t} \pi'_*(\mathcal{A}) q_3 \xrightarrow{h_2} \pi'_*(\mathcal{A}) q_4.$$

Hence, there exist $k_1, k_2 \in H$ with $h_1 = \pi'(k_1)$, $h_2 = \pi'(k_2)$, and

$$q_1 \xrightarrow{k_1} \mathcal{A} q_2 \xrightarrow{t} \mathcal{A} q_3 \xrightarrow{k_2} \mathcal{A} q_4.$$

Let $a \in A$. Since \mathcal{A} is \sim -saturated, there are states $p_2, p_3 \in \mathcal{Q}$ such that

$$q_1 \xrightarrow{k_1 a^{-1}} \mathcal{A} p_2 \xrightarrow{t} \mathcal{A} p_3 \xrightarrow{\varphi(a)k_2} \mathcal{A} q_4$$

and hence

$$q_1 \xrightarrow{\pi'(k_1 a^{-1})} \pi'_*(\mathcal{A}) p_2 \xrightarrow{t} \pi'_*(\mathcal{A}) p_3 \xrightarrow{\pi'(\varphi(a)k_2)} \pi'_*(\mathcal{A}) q_4.$$

Since $\pi'(c) = c$ for all $c \in A \cup B$, we obtain

$$q_1 \xrightarrow{h_1 a^{-1}}_{\pi'_*(\mathcal{A})} p_2 \xrightarrow{t}_{\pi'_*(\mathcal{A})} p_3 \xrightarrow{\varphi(a)h_2}_{\pi'_*(\mathcal{A})} q_4.$$

An analogous argument for t^{-1} shows that $\pi'_*(\mathcal{A})$ is \sim' -saturated. \square

4.4. Normalization of automata

The following proposition is crucial for our further considerations.

Proposition 22. *From a given partitioned and \approx -compatible fta \mathcal{A} with labelling set $\mathcal{F} \subseteq \mathcal{P}(H)$ one can effectively construct a partitioned, \approx -compatible, \sim -saturated, and unitary fta \mathcal{B} with labelling set $\{cF c' \mid F \in \mathcal{F}, c, c' \in A \cup B\}$ such that $[\mathbf{L}(\mathcal{A})]_{\approx} \cap \text{Red}(H, t) = \mathbf{L}(\mathcal{B}) \cap \text{Red}(H, t)$.*

Proof. Let $\mathcal{A} = \langle \mathcal{L}, \mathbf{Q}, \tau, \delta, \mathbf{l}, \mathbf{T} \rangle$ be some partitioned and \approx -compatible fta with labeling set $\mathcal{F} \subseteq \mathcal{P}(H)$. Thus, we have

$$[\mathbf{L}(\mathcal{A})]_{\approx} = [\mathbf{L}(\mathcal{A}) \cap \text{Red}(H, t)]_{\approx}. \quad (37)$$

Since \mathcal{A} is partitioned, Lemma 7 implies that we can replace the set of transitions δ by the set

$$\{(p, L, q) \in \delta \mid \tau(p) \neq \tau(q)\} \cup \{(q, 1, q) \mid q \in \mathbf{Q}\}$$

and preserve the recognized language. The second term of the above union ensures that property (23) still holds. Such a modification might modify the mappings $\mu_{\mathcal{A}, 0}$ and $\mu_{\mathcal{A}}$, but for every sequence s it does not modify the relation $\mu_{\mathcal{A}}(s) \cap \tau^{-1}(1, H) \times \tau^{-1}(1, 1)$, i.e., after this modification, \mathcal{A} is still a partitioned and \approx -compatible fta. Hence, w.l.o.g. we assume that

$$\forall (p, L, q) \in \delta : \tau(p) \neq \tau(q) \vee (p = q \wedge L = \{1\}). \quad (38)$$

Note that property (38) implies that \mathcal{A} is unitary according to Definition 11.

We define the fta $\mathcal{B} = \langle \mathcal{L}_{\mathcal{B}}, \mathbf{Q}_{\mathcal{B}}, \tau_{\mathcal{B}}, \delta_{\mathcal{B}}, \mathbf{l}, \mathbf{T} \rangle$ in the following way: Let us denote for every $q \in \mathbf{Q}$ by $\gamma(q)$ the subgroup


$$\gamma(q) = \pi_1(\tau(q)) \in \{1, A, B\}$$

where π_1 denotes the projection onto the first component of a type. Since \mathcal{A} is partitioned, if $(q, t, r) \in \delta$ (resp. $(q, t^{-1}, r) \in \delta$) then $\gamma(q) = A$ (resp. $\gamma(r) = B$) and $\gamma(q) = B$ (resp. $\gamma(r) = A$). We set

$$\mathbf{Q}_{\mathcal{B}} = \{(q, g) \in \mathbf{Q} \times (A \cup B) \mid g \in \gamma(q)\}.$$

We shall identify $(q, 1)$ with q and denote (q, g) by qg in the sequel. Note that the set of initial (resp. terminal) states of \mathcal{B} is the same as for \mathcal{A} . The typing map $\tau_{\mathcal{B}}$ is inherited from \mathcal{A} :

$$\tau_{\mathcal{B}}(qg) = \tau(q)$$


 Fig. 4. Some subgraphs of \mathcal{B}

The set $\delta_{\mathcal{B}}$ consists of all the following transitions, where $q, r \in \mathcal{Q}$:

$$(qg, g', q(gg')) \text{ for } g, g' \in \gamma(q) \quad (39)$$

$$(qg, g^{-1}Lg', rg') \text{ for } g \in \gamma(q), g' \in \gamma(r), L \subseteq H, (q, L, r) \in \delta \quad (40)$$

$$(qa, t, r\varphi(a)) \text{ for } a \in A, (q, t, r) \in \delta \quad (41)$$

$$(qb, t^{-1}, r\varphi^{-1}(b)) \text{ for } b \in B, (q, t^{-1}, r) \in \delta \quad (42)$$

For all $(q_1, L, r_1), (q, t, r), (\bar{q}, t^{-1}, \bar{r}) \in \delta$, $g \in \gamma(q_1)$, $g' \in \gamma(r_1)$, $h \in L$, $a \in A$, and $b \in B$, the automaton \mathcal{B} contains the three basic subgraphs exhibited in Figure 4. We cut the proof that \mathcal{B} has the required properties into five claims.

Claim 23. $[\mathcal{L}(\mathcal{A})]_{\approx} \cap \text{Red}(H, t) = [\mathcal{L}(\mathcal{A}) \cap \text{Red}(H, t)]_{\approx}$

By intersecting both sides of (37) with $\text{Red}(H, t)$, we obtain

$$[\mathcal{L}(\mathcal{A})]_{\approx} \cap \text{Red}(H, t) = [\mathcal{L}(\mathcal{A}) \cap \text{Red}(H, t)]_{\approx} \cap \text{Red}(H, t).$$

But by Lemma 5, the right-hand side above equals $[\mathcal{L}(\mathcal{A}) \cap \text{Red}(H, t)]_{\approx}$, which proves Claim 23.

Claim 24. $\mathcal{L}(\mathcal{A}) \subseteq \mathcal{L}(\mathcal{B})$

Let us notice that every $(q, L, r) \in \delta$ has a copy in $\delta_{\mathcal{B}}$ under the form of (q_1, L, r_1) , since whatever $\gamma(q)$ and $\gamma(r)$ are, $1 \in \gamma(q) \cap \gamma(r)$. Thus, up to the identification of every state $q \in \mathcal{Q}$ with the state $q_1 \in \mathcal{Q}_{\mathcal{B}}$, $\delta \subseteq \delta_{\mathcal{B}}$. Since moreover $\mathcal{I} = \mathcal{I}_{\mathcal{B}}$ and $\mathcal{T} = \mathcal{T}_{\mathcal{B}}$, we obtain Claim 24.

Claim 25. *The fta \mathcal{B} is \sim -saturated and hence, $\mathcal{L}(\mathcal{B}) = [\mathcal{L}(\mathcal{B})]_{\approx}$.*

Let us assume that

$$q_0a_0 \xrightarrow{\mathcal{B}}_h q_1a_1 \xrightarrow{\mathcal{B}}_t q_2a_2 \xrightarrow{\mathcal{B}}_{h'} q_3a_3,$$

where $q_0, \dots, q_3 \in \mathcal{Q}$, $a_i \in \gamma(q_i)$, and $h, h' \in H$. Thus, $(q_1, t, q_2) \in \delta$ and $a_2 = \varphi(a_1)$. Moreover, $\gamma(q_1) = A$, since a t -transition is enabled in q_1a_1 . The transition $q_0a_0 \xrightarrow{\mathcal{B}}_h q_1a_1$ is either of type (39) or (40). In both cases, we can derive for every $a \in A$ the transition

$$q_0a_0 \xrightarrow{\mathcal{B}}_{ha^{-1}} q_1(a_1a^{-1}). \quad (43)$$

Analogously, we get

$$q_2\varphi(a_1a^{-1}) \xrightarrow{\mathcal{B}}_{\varphi(a)h'} q_3a_3.$$

Finally, $q_1 a_1 \xrightarrow{t}_{\mathcal{B}} q_2 \varphi(a_1)$ and (41) implies

$$q_1(a_1 a^{-1}) \xrightarrow{t}_{\mathcal{B}} q_2 \varphi(a_1 a^{-1}).$$

Altogether, it follows

$$q_0 a_0 \xrightarrow{(ha^{-1})t(\varphi(a)h')}_{\mathcal{B}} q_3 a_3. \quad (45)$$

Similarly, if $q_0 a_0 \xrightarrow{ht^{-1}h'}_{\mathcal{B}} q_3 a_3$ and $b \in B$, then

$$q_0 a_0 \xrightarrow{(hb^{-1})t^{-1}(\varphi^{-1}(b)h')}_{\mathcal{B}} q_3 a_3.$$

This proves Claim 25.

Claim 26. $L(\mathcal{B}) \subseteq [L(\mathcal{A})]_{\sim}$

Let $s \in H * \{t, t^{-1}\}^*$, $q, r \in \mathbf{Q}$, $g \in \gamma(q)$, and $h \in \gamma(r)$ such that

$$qg \xrightarrow{s}_{\mathcal{B}} rh. \quad (46)$$

By gluing squares from Figure 4 along their vertical edges, one can prove by induction on the integer n of the decomposition (15) that, under the hypothesis (46), there exists $s' \sim gsh^{-1}$ such that

$$q \xrightarrow{s'}_{\mathcal{A}} r. \quad (47)$$

If $q \in \mathbf{l}$ and $r \in \mathbf{T}$, then the type-conditions (30) imply $\gamma(q) = \gamma(r) = 1$, i.e., $g = h = 1$ and $s \in L(\mathcal{B})$. Then (47) says that

$$\exists s' : s' \sim s \wedge s' \in L(\mathcal{A}),$$

which proves Claim 26.

Claim 27. $L(\mathcal{B}) \subseteq [L(\mathcal{B}) \cap \text{Red}(H, t)]_{\approx}$

Claim 27 can be deduced as follows:

$$\begin{aligned} L(\mathcal{B}) &\subseteq [L(\mathcal{A})]_{\sim} && \text{(by Claim 26)} \\ &\subseteq [L(\mathcal{A})]_{\approx} && \text{(since } \sim \subseteq \approx \text{)} \\ &= [L(\mathcal{A}) \cap \text{Red}(H, t)]_{\approx} && \text{(since } \mathcal{A} \text{ is } \approx\text{-compatible)} \\ &\subseteq [L(\mathcal{B}) \cap \text{Red}(H, t)]_{\approx} && \text{(by Claim 24)} \end{aligned}$$

Let us now use Claim 23–27 in order to show point Proposition 22: First, we have

$$\begin{aligned} [L(\mathcal{A})]_{\approx} \cap \text{Red}(H, t) &= [L(\mathcal{A}) \cap \text{Red}(H, t)]_{\sim} && \text{(by Claim 23)} \\ &\subseteq [L(\mathcal{A})]_{\sim} \\ &\subseteq [L(\mathcal{B})]_{\sim} && \text{(by Claim 24)} \\ &= L(\mathcal{B}) && \text{(by Claim 25),} \end{aligned}$$

and thus

$$[L(\mathcal{A})]_{\approx} \cap \text{Red}(H, t) \subseteq L(\mathcal{B}) \cap \text{Red}(H, t).$$

On the other hand, we have

$$\begin{aligned} \mathbf{L}(\mathcal{B}) \cap \text{Red}(H, t) &\subseteq [\mathbf{L}(\mathcal{A})]_{\sim} \cap \text{Red}(H, t) \quad (\text{by Claim 26}) \\ &\subseteq [\mathbf{L}(\mathcal{A})]_{\approx} \cap \text{Red}(H, t) \quad (\text{since } \sim \subseteq \approx). \end{aligned}$$

Hence, we obtain

$$[\mathbf{L}(\mathcal{A})]_{\approx} \cap \text{Red}(H, t) = \mathbf{L}(\mathcal{B}) \cap \text{Red}(H, t),$$

as was required. Moreover:

- \mathcal{B} is partitioned, since \mathcal{A} is partitioned.
- By Claim 27, \mathcal{B} is \approx -compatible.
- By Claim 25, \mathcal{B} is \sim -saturated.
- \mathcal{B} is unitary: If $g' = 1$ in (39), then we have $qg = qgg'$. If $1 \in g^{-1}Lg'$ and $(q, L, r) \in \delta$ in (40), then property (38) guarantees that $\tau(q) \neq \tau(r)$ or $(q = r \text{ and } L = \{1\})$. If $\tau(q) \neq \tau(r)$ then $\tau_{\mathcal{B}}(qg) \neq \tau_{\mathcal{B}}(rg')$. If $q = r$ and $L = \{1\}$, then $g = g'$ and hence $qg = rg'$. Hence, no \mathcal{B} -transition of type (39) or (40) has the form (qg, L, rg') with $\tau_{\mathcal{B}}(qg) = \tau_{\mathcal{B}}(rg')$, $1 \in L$, and $qg \neq rg'$.

Proposition 22 is thus established. \square

Recall the definition of a deterministic (Definition 12) and complete (Definition 13) fta.

Proposition 28. *From a given partitioned, \approx -compatible, and \sim -saturated fta \mathcal{B} with labelling set $\mathcal{F} \subseteq \mathcal{P}(H)$ one can effectively construct a partitioned, \sim -saturated, deterministic, and complete fta \mathcal{C} with labelling set $\text{Bool}(\mathcal{F})$ such that $\mathbf{L}(\mathcal{B}) \cap \text{Red}(H, t) = \mathbf{L}(\mathcal{C})$.*

Proof. The proof is essentially a variant of the standard power set construction for determinizing ordinary finite state automata over words. Let $\mathcal{B} = \langle \mathcal{L}_{\mathcal{B}}, \mathbf{Q}_{\mathcal{B}}, \tau_{\mathcal{B}}, \delta_{\mathcal{B}}, \mathbf{l}_{\mathcal{B}}, \mathbf{T}_{\mathcal{B}} \rangle$ be a partitioned, \approx -compatible, and \sim -saturated fta. We first define a partitioned fta

$$\mathcal{D} = \langle \mathcal{L}_{\mathcal{D}}, \mathbf{Q}_{\mathcal{D}}, \tau_{\mathcal{D}}, \delta_{\mathcal{D}}, \mathbf{l}_{\mathcal{D}}, \mathbf{T}_{\mathcal{D}} \rangle$$

in the following way: Let

$$\mathcal{L}' = (\mathcal{L}_{\mathcal{B}} \cap \mathcal{P}(H)) \cup \{\{c\} \mid c \in A \cup B\} \quad (49)$$

and

$$\mathcal{L}_{\mathcal{D}} = \{\{t\}, \{t^{-1}\}\} \cup \left\{ \bigcap_{L \in \mathcal{K}} L \cap \bigcap_{L \in \mathcal{L}' \setminus \mathcal{K}} H \setminus L \mid \mathcal{K} \subseteq \mathcal{L}' \right\} \setminus \{\emptyset\}.$$

Thus, $\mathcal{L}_{\mathcal{D}} \cap \mathcal{P}(H)$ is the set of atoms of the boolean algebra generated by \mathcal{L}' . Let

$$\mathbf{Q}_{\mathcal{D}} = \{(P, \Theta) \mid P \subseteq \mathbf{Q}_{\mathcal{B}}, \Theta \in \mathcal{T}_6, P \subseteq \tau_{\mathcal{B}}^{-1}(\Theta)\}.$$

The type function is the projection onto the second component:

$$\tau_{\mathcal{D}}(P, \Theta) = \Theta.$$

The set of transitions $\delta_{\mathcal{D}}$ consists of all triples $((P, \Theta), K, (P', \Theta')) \in \mathcal{Q}_{\mathcal{D}} \times \mathcal{L}_{\mathcal{D}} \times \mathcal{Q}_{\mathcal{D}}$ such that

$$\begin{aligned} P' &= \{p' \in \mathcal{Q}_{\mathcal{B}} \mid \exists p \in P \exists L : (p, L, p') \in \delta_{\mathcal{B}}, K \subseteq L\} \cap \tau_{\mathcal{B}}^{-1}(\Theta') \\ &\text{and } \{\Theta\} \times K \times \{\Theta'\} \subseteq \widehat{\mathcal{E}}_6, \end{aligned}$$

where \mathcal{E}_6 is the edge relation of the graph \mathcal{G}_6 and $\widehat{\mathcal{E}}_6$ is defined by (24). Note that $P' = \emptyset$ is allowed. Finally, let

$$\mathsf{l}_{\mathcal{D}} = \{(\mathsf{l}_{\mathcal{B}}, (1, H))\} \text{ and } \mathsf{T}_{\mathcal{D}} = \{(P, (1, 1)) \in \mathcal{Q}_{\mathcal{D}} \mid P \cap \mathsf{T}_{\mathcal{B}} \neq \emptyset\}.$$

The definition of $\delta_{\mathcal{D}}$ implies that \mathcal{D} is partitioned and complete. Moreover, the fta \mathcal{D} is deterministic: if we have two transitions $((P, \Theta), K_i, (P_i, \Theta')) \in \delta_{\mathcal{D}}$ ($i \in \{1, 2\}$) with $K_1 \cap K_2 \neq \emptyset$ then we must have $K_1 = K_2$ and hence $P_1 = P_2$.

Remark 29. For every $P \subseteq \mathcal{Q}_{\mathcal{B}}$ fulfilling $\emptyset \subsetneq P \subseteq \tau^{-1}(\Theta)$, (P, Θ) is a state of \mathcal{D} and no pair (P, Θ') with $\Theta \neq \Theta'$ is a state of \mathcal{D} . But (\emptyset, Θ) is a state of \mathcal{D} for every type Θ . This is in some sense the price to pay for the completeness of \mathcal{D} .

Recall the definition of the mapping $\mu_{\mathcal{B},0}$ from (25).

Claim 30. *Let $(P, \Theta), (P', \Theta') \in \mathcal{Q}_{\mathcal{D}}$ and $s \in H \cup \{t, t^{-1}\}$. Then*

$$((P, \Theta), (P', \Theta')) \in \mu_{\mathcal{D},0}(s) \iff P' = (P \odot \mu_{\mathcal{B},0}(s)) \cap \tau_{\mathcal{B}}^{-1}(\Theta') \wedge (\Theta, s, \Theta') \in \widehat{\mathcal{E}}_6.$$

Remark 31. If $P' = (P \odot \mu_{\mathcal{B},0}(s)) \cap \tau_{\mathcal{B}}^{-1}(\Theta') \neq \emptyset$, then the condition $(\Theta, s, \Theta') \in \widehat{\mathcal{E}}_6$ follows automatically, since the fta \mathcal{B} is partitioned. But since $P' = \emptyset$ is possible, we have to add the condition $(\Theta, s, \Theta') \in \widehat{\mathcal{E}}_6$ explicitly.

In order to prove Claim 30, we distinguish three cases.

Case 1: $s = t$. Note that $((A, T), t, (B, H))$ is the only t -labelled edge of \mathcal{G}_6 . We get:

$$\begin{aligned} &((P, \Theta), (P', \Theta')) \in \mu_{\mathcal{D},0}(t) \\ &\iff ((P, \Theta), t, (P', \Theta')) \in \delta_{\mathcal{D}} \\ &\iff P' = P \odot \mu_{\mathcal{B},0}(t) \cap \tau_{\mathcal{B}}^{-1}(\Theta') \wedge \Theta = (A, T) \wedge \Theta' = (B, H). \end{aligned}$$

Case 2: $s = t^{-1}$. Same kind of arguments as for $s = t$.

Case 3: $s = h \in H$. Let $K(h)$ be the unique atom of $\mathcal{L}_{\mathcal{D}} \cap \mathcal{P}(H)$ such that $h \in K(h)$. For every transition $(p, L, q) \in \delta_{\mathcal{B}}$, one has

$$h \in L \iff K(h) \subseteq L.$$

It follows that

$$\begin{aligned}
 ((P, \Theta), (P', \Theta')) \in \mu_{\mathcal{D},0}(h) &\Leftrightarrow ((P, \Theta), K(h), (P', \Theta')) \in \delta_{\mathcal{D}} \\
 &\Leftrightarrow P' = (P \odot \mu_{\mathcal{B},0}(h)) \cap \tau_{\mathcal{B}}^{-1}(\Theta') \text{ and} \\
 &\quad \{\Theta\} \times K(h) \times \{\Theta'\} \subseteq \widehat{\mathcal{E}}_6 \\
 &\Leftrightarrow P' = (P \odot \mu_{\mathcal{B},0}(h)) \cap \tau_{\mathcal{B}}^{-1}(\Theta') \text{ and} \\
 &\quad (\Theta, h, \Theta') \in \widehat{\mathcal{E}}_6.
 \end{aligned}$$

For the last equivalence note that $\{\Theta\} \times K(h) \times \{\Theta'\} \subseteq \widehat{\mathcal{E}}_6$ if and only if $(\Theta, h, \Theta') \in \widehat{\mathcal{E}}_6$. For this, it is important that if $h \in A \cup B$ then $K(h) = \{h\}$; note that every singleton $\{c\}$ with $c \in A \cup B$ belongs to \mathcal{L}' in (49).

Claim 32. *Let $(P, \Theta), (P', \Theta') \in \mathbf{Q}_{\mathcal{D}}$ and $s \in H * \{t, t^{-1}\}^*$. Then*

$$((P, \Theta), (P', \Theta')) \in \mu_{\mathcal{D}}(s) \Leftrightarrow P' = (P \odot \mu_{\mathcal{B}}(s)) \cap \tau_{\mathcal{B}}^{-1}(\Theta') \wedge (\Theta, \Theta') \in \mu_{\mathcal{G}_6}(s).$$

Let us prove this claim by induction over the integer n appearing in decomposition (15) of the t -sequence s .

Base case: $n = 0$. For $s = h \in H$ the assertion of Claim 32 is equivalent with that of Claim 30, since $\mu_{\mathcal{A}}(h) = \mu_{\mathcal{A},0}(h)$ for every fta \mathcal{A} .

Induction step: Let s be of the form (15) with $n \geq 1$. We suppose that $\alpha_n = 1$ and let

$$s_1 = h_0 t^{\alpha_1} h_1 \cdots t^{\alpha_i} h_i \cdots t^{\alpha_{n-1}} h_{n-1} \text{ and } s = s_1 t h_n$$

(where $s_1 = h_0$ if $n = 1$). Let us first assume that

$$((P, \Theta), (P', \Theta')) \in \mu_{\mathcal{D}}(s). \quad (50)$$

The definition of $\delta_{\mathcal{D}}$ implies $(\Theta, \Theta') \in \mu_{\mathcal{G}_6}(s)$. Moreover, the structure of the graph \mathcal{G}_6 implies that there exist $P_1 \subseteq \tau_{\mathcal{B}}^{-1}(A, T)$ and $P_2 \subseteq \tau_{\mathcal{B}}^{-1}(B, H)$ with

$$((P, \Theta), (P_1, (A, T))) \in \mu_{\mathcal{D}}(s_1), \quad (51)$$

$$((P_1, (A, T)), (P_2, (B, H))) \in \mu_{\mathcal{D},0}(t), \text{ and} \quad (52)$$

$$((P_2, (B, H)), (P', \Theta')) \in \mu_{\mathcal{D},0}(h_n). \quad (53)$$

As Claim 32 is fulfilled by s_1 (by the induction hypothesis) and also by sequences in $H \cup \{t, t^{-1}\}$ (by Claim 30), equations (51)–(53) imply:

$$(P \odot \mu_{\mathcal{B}}(s_1)) \cap \tau_{\mathcal{B}}^{-1}(A, T) = P_1 \quad (54)$$

$$(P_1 \odot \mu_{\mathcal{B},0}(t)) \cap \tau_{\mathcal{B}}^{-1}(B, H) = P_2 \quad (55)$$

$$(P_2 \odot \mu_{\mathcal{B},0}(h_n)) \cap \tau_{\mathcal{B}}^{-1}(\Theta') = P' \quad (56)$$

Omitting the type intersections in (54) and (55), we obtain

$$(P \odot \mu_{\mathcal{B}}(s_1 t h_n)) \cap \tau_{\mathcal{B}}^{-1}(\Theta) \supseteq P'. \quad (57)$$

Conversely, by the structure of \mathcal{G}_6 , every path in the partitioned fta \mathcal{B} from a state $p \in P$ to a state $q' \in \tau_{\mathcal{B}}^{-1}(\Theta')$, reading s_1th_n , must decompose into

$$p \xrightarrow{s_1}_{\mathcal{B}} p_1 \xrightarrow{t}_{\mathcal{B}} p_2 \xrightarrow{h_n}_{\mathcal{B}} q'$$

with $\tau_{\mathcal{B}}(p_1) = (A, T)$ and $\tau_{\mathcal{B}}(p_2) = (B, H)$. Hence, (54)–(56) imply

$$(P \odot \mu_{\mathcal{B}}(s_1th_n)) \cap \tau_{\mathcal{B}}^{-1}(\Theta') \subseteq P'. \quad (58)$$

From (57) and (58) we obtain $P \odot \mu_{\mathcal{B}}(s_1th_n) \cap \tau_{\mathcal{B}}^{-1}(\Theta') = P'$. This proves implication \Rightarrow from Claim 32.

Now, suppose conversely that $P \odot \mu_{\mathcal{B}}(s_1th_n) \cap \tau_{\mathcal{B}}^{-1}(\Theta') = P'$ and $(\Theta, \Theta') \in \mu_{\mathcal{G}_6}(s)$. By the same kind of arguments as above, there must exist $P_1 \subseteq \tau_{\mathcal{B}}^{-1}(A, T)$ and $P_2 \subseteq \tau_{\mathcal{B}}^{-1}(B, H)$ fulfilling equations (54)–(56). Moreover, from $(\Theta, \Theta') \in \mu_{\mathcal{G}_6}(s)$ we get $(\Theta, (A, T)) \in \mu_{\mathcal{G}_6}(s_1)$ and $((B, H), \Theta') \in \mu_{\mathcal{G}_6}(h_n)$. By the induction hypothesis and Claim 30 this leads to (51)–(53), and hence to (50).

The induction step in the case where $\alpha_n = -1$ can be treated analogously. This proves Claim 32.

From Claim 32 and the fact that the fta \mathcal{B} and \mathcal{G}_6 are \sim -saturated, it follows immediately that also \mathcal{D} is \sim -saturated. Moreover,

$$L(\mathcal{D}) = L(\mathcal{B}).$$

Let us finally take for \mathcal{C} the product automaton (see Definition 14) $\mathcal{C} = \mathcal{D} \times \mathcal{R}_6$, where \mathcal{R}_6 is the partitioned fta from Figure 3. The partitioned fta \mathcal{C} has the following properties:

- $L(\mathcal{C}) = L(\mathcal{D}) \cap \text{Red}(H, t) = L(\mathcal{B}) \cap \text{Red}(H, t)$ by Lemma 15.
- \mathcal{C} is deterministic and complete by Lemma 18 and the fact that \mathcal{R}_6 and \mathcal{D} are both deterministic and complete.
- \mathcal{C} is \sim -saturated by Lemma 17 and the fact that \mathcal{R}_6 and \mathcal{D} are \sim -saturated.

Hence, \mathcal{C} fulfils Proposition 28. \square

Recall that $\pi : (X \cup \{t, t^{-1}\})^* \rightarrow G = \langle H, t; t^{-1}at = \varphi(a)(a \in A) \rangle$ denotes the canonical morphism from (8), where X is a generating set for H with $A \cup B \subseteq X$.

Proposition 33. *Let $K \subseteq G$. The following are equivalent:*

- (1) K is a rational subset of G .
- (2) There exists some finite automaton \mathcal{A} over the free monoid $(X \cup \{t, t^{-1}\})^*$ such that $K = \pi(L(\mathcal{A}))$.
- (3) $\pi_G^{-1}(K) \cap \text{Red}(H, t) = L(\mathcal{B}) \cap \text{Red}(H, t)$ for some partitioned, \approx -compatible, and \sim -saturated fta \mathcal{B} with labelling set $\text{Rat}(H)$.

Moreover,

- (a) in point (3), \mathcal{B} can always be chosen unitary (see Definition 11), and

(b) in point (3), if the membership problem for languages in $\text{Rat}(H)$ is decidable, then \mathcal{B} can be computed effectively from \mathcal{A} in point (2).

Proof. The equivalence of (1) and (2) is trivial.

(2) \Rightarrow (3): We cut the proof of this implication into three claims. Let us start with a finite automaton

$$\mathcal{A} = \langle \mathcal{X}, \mathbf{Q}, \delta, \mathbf{l}, \mathbf{T} \rangle$$

over the free monoid $(X \cup \{t, t^{-1}\})^*$ such that

$$K = \pi(\mathbf{L}(\mathcal{A})).$$

Let us recall that ST is the semi-Thue system consisting of all the rules in (16) and (17), where $u \in X^*$ and $\varphi([u]_{\equiv}) \in B$ and $\varphi^{-1}([u]_{\equiv}) \in A$ denote the corresponding letters in X (we choose such a letter for every element of the finite set $A \cup B$).

For a set E , a binary relation $\rightsquigarrow \subseteq E \times E$, and a subset $R \subseteq E$, let $R \rightsquigarrow = \{e \in E \mid \exists r \in R : r \rightsquigarrow e\}$.

Claim 34. *There exists a finite automaton \mathcal{A}_1 over the free monoid $(X \cup \{t, t^{-1}\})^*$, such that*

$$\mathbf{L}(\mathcal{A}_1) = \mathbf{L}(\mathcal{A}) \rightarrow_{\text{ST}}^*.$$

Moreover, if the membership problem is decidable for rational subsets of H , then \mathcal{A}_1 can be constructed effectively from \mathcal{A} .

For the proof of Claim 34, notice that ST is *monadic*, i.e., every right-hand side of a rule has length one or zero. It is known that for every monadic semi-Thue system S over an alphabet Y and every rational subset $R \subseteq Y^*$, $R \rightarrow_S^* \in \text{Rat}(Y^*)$, see [BJW82, S en94]. Taking $R = \mathbf{L}(\mathcal{A})$ and $S = \text{ST}$ in this result, we obtain the first statement of Claim 34. In order to check the effectiveness statement, let us consider the following algorithm:

```

1   $\delta_1 := \delta$ ;
2  while there are  $p, q, q', r \in \mathbf{Q}, u \in X^*, a \in A, b \in B$  such that
    case 1:  $p \xrightarrow{t^{-1}}_{\delta_1} q \xrightarrow{u}_{\delta_1} q' \xrightarrow{t}_{\delta_1} r, [u]_{\equiv} = a$ , and not  $p \xrightarrow{\varphi(a)}_{\delta_1} r$ 
    or
    case 2:  $p \xrightarrow{t}_{\delta_1} q \xrightarrow{u}_{\delta_1} q' \xrightarrow{t^{-1}}_{\delta_1} r, [u]_{\equiv} = b$ , and not  $p \xrightarrow{\varphi^{-1}(b)}_{\delta_1} r$ 
    do
3      in case 1:  $\delta_1 := \delta_1 \cup \{(p, \varphi(a), r)\}$ 
4      in case 2:  $\delta_1 := \delta_1 \cup \{(p, \varphi^{-1}(b), r)\}$ 
5  od
    
```

This algorithm constructs from the finite automaton \mathcal{A} a finite automaton

$$\mathcal{A}_1 = \langle \mathcal{X} \cup A \cup B, \mathbf{Q}, \delta_1, \mathbf{l}, \mathbf{T} \rangle \quad (59)$$

satisfying the first statement of Claim 34. In order to make line 2 above effective, it is enough that we can decide for every $a \in A$ (resp. $b \in B$) and every $q, q' \in \mathbb{Q}$, whether $a \in \pi_t(\mathcal{L}(\mathcal{A}[X, q, q']))$ (resp. $b \in \pi_t(\mathcal{L}(\mathcal{A}[X, q, q'])))$, where $\pi_t : (X \cup \{t, t^{-1}\})^* \rightarrow H * \{t, t^{-1}\}^*$ is the canonical morphism from (10) and the automaton $\mathcal{A}[X, q, q']$ is defined by (5). But this is an instance of the membership problem for rational subsets of H . This proves the effectiveness statement in Claim 34.

Claim 35. *There exists some \approx -compatible fta \mathcal{A}_2 over $H * \{t, t^{-1}\}^*$ with labelling set $\text{Rat}(H)$ such that*

$$\mathcal{L}(\mathcal{A}_2) = \pi_t(\mathcal{L}(\mathcal{A}_1)).$$

Moreover, \mathcal{A}_2 can be constructed effectively from \mathcal{A}_1 .

Recall the finite automaton \mathcal{A}_1 from (59). We set

$$\mathcal{A}_2 = \langle \mathcal{L}, \mathbb{Q}, \delta_2, \mathbb{1}, \mathbb{T} \rangle$$

where

$$\begin{aligned} \delta_2 = \{ & (q, t^\alpha, r) \mid \alpha \in \{1, -1\}, (q, t^\alpha, r) \in \delta_1 \} \cup \\ & \{(q, \pi_t(\mathcal{L}(\mathcal{A}_1[X, q, r])), r) \mid q, r \in \mathbb{Q}\} \end{aligned}$$

(the label set \mathcal{L} is given implicitly by δ_2). In some sense the transitions of δ_2 are collecting into the rational labels $\pi_t(\mathcal{L}(\mathcal{A}_1[X, q, r])) \in \text{Rat}(H)$ all possible finite paths with labels in X^* , while the transitions labelled by t or t^{-1} are just copied. Clearly, $\mathcal{L}(\mathcal{A}_2) = \pi_t(\mathcal{L}(\mathcal{A}_1))$ and \mathcal{A}_2 can be constructed effectively from \mathcal{A}_1 . So it remains to show that \mathcal{A}_2 is \approx -compatible.

Recall that $\pi_t(\text{Irr}(\text{ST})) = \text{Red}(H, t)$. Since, $\mathcal{L}(\mathcal{A}_1) = \mathcal{L}(\mathcal{A}) \rightarrow_{\text{ST}}^*$, the language $\mathcal{L}(\mathcal{A}_1)$ is closed under the terminating rewrite relation $\rightarrow_{\text{ST}} \subseteq \equiv$. It follows

$$\mathcal{L}(\mathcal{A}_1) \subseteq [\mathcal{L}(\mathcal{A}_1) \cap \text{Irr}(\text{ST})]_{\equiv}.$$

By applying the morphism π_t to this inclusion, we obtain

$$\begin{aligned} \mathcal{L}(\mathcal{A}_2) &= \pi_t(\mathcal{L}(\mathcal{A}_1)) \\ &\subseteq \pi_t([\mathcal{L}(\mathcal{A}_1) \cap \text{Irr}(\text{ST})]_{\equiv}) \\ &= [\pi_t(\mathcal{L}(\mathcal{A}_1) \cap \text{Irr}(\text{ST}))]_{\approx} \\ &\subseteq [\pi_t(\mathcal{L}(\mathcal{A}_1)) \cap \pi_t(\text{Irr}(\text{ST}))]_{\approx} \\ &= [\mathcal{L}(\mathcal{A}_2) \cap \text{Red}(H, t)]_{\approx}, \end{aligned}$$

which proves that \mathcal{A}_2 is \approx -compatible.

Claim 36. *There exists some partitioned and \approx -compatible fta \mathcal{A}_3 over $H * \{t, t^{-1}\}^*$ with labelling set $\text{Rat}(H)$ such that*

$$\mathcal{L}(\mathcal{A}_3) = \mathcal{L}(\mathcal{A}_2).$$

Moreover, \mathcal{A}_3 can be constructed effectively from \mathcal{A}_2 .

Just set $\mathcal{A}_3 = \mathcal{A}_2 \times \mathcal{G}_6$. By Lemma 16, we have $L(\mathcal{A}_3) = L(\mathcal{A}_2)$. Hence, since \mathcal{A}_2 is \approx -compatible, also \mathcal{A}_3 is \approx -compatible.

By Proposition 22 there exists a partitioned, \approx -compatible, \sim -saturated, and unitary fta \mathcal{B} with labelling set $\text{Rat}(H)$ such that

$$L(\mathcal{B}) \cap \text{Red}(H, t) = [L(\mathcal{A}_3)]_{\approx} \cap \text{Red}(H, t).$$

Let us summarize the connections between the rational set $K \subseteq G$ and the successive automata $\mathcal{A}, \mathcal{A}_1, \mathcal{A}_2, \mathcal{A}_3$, and \mathcal{B} :

$$K = \pi(L(\mathcal{A})) \subseteq G \quad (60)$$

$$L(\mathcal{A}_1) = L(\mathcal{A}) \xrightarrow{\ast} \text{S}_T \subseteq (X \cup \{t, t^{-1}\})^* \quad (61)$$

$$L(\mathcal{A}_2) = \pi_t(L(\mathcal{A}_1)) \subseteq H * \{t, t^{-1}\}^* \quad (62)$$

$$L(\mathcal{A}_3) = L(\mathcal{A}_2) \subseteq H * \{t, t^{-1}\}^* \quad (63)$$

$$L(\mathcal{B}) \cap \text{Red}(H, t) = [L(\mathcal{A}_3)]_{\approx} \cap \text{Red}(H, t) \subseteq H * \{t, t^{-1}\}^* \quad (64)$$

Since $\pi = \pi_t \circ \pi_G$ (9), equation (60) implies

$$\pi_G^{-1}(K) = \pi_G^{-1}(\pi_G(\pi_t(L(\mathcal{A}))) = [\pi_t(L(\mathcal{A}))]_{\approx}.$$

By (61), $[L(\mathcal{A})]_{\equiv} = [L(\mathcal{A}_1)]_{\equiv}$, hence $[\pi_t(L(\mathcal{A}))]_{\approx} = [\pi_t(L(\mathcal{A}_1))]_{\approx}$ and thus

$$\pi_G^{-1}(K) = [\pi_t(L(\mathcal{A}_1))]_{\approx}.$$

Using (62) and (63) we get

$$\pi_G^{-1}(K) = [L(\mathcal{A}_2)]_{\approx} = [L(\mathcal{A}_3)]_{\approx}.$$

Finally, by (64) we have

$$\pi_G^{-1}(K) \cap \text{Red}(H, t) = L(\mathcal{B}) \cap \text{Red}(H, t).$$

We have obtained point (3) of Proposition 33.

(3) \Rightarrow (1): Let \mathcal{B} be some partitioned, \approx -compatible, and \sim -saturated fta with labelling set $\text{Rat}(H)$ such that

$$\pi_G^{-1}(K) \cap \text{Red}(H, t) = L(\mathcal{B}) \cap \text{Red}(H, t). \quad (65)$$

Let $\mathcal{L}_{\mathcal{B}} \subseteq \text{Rat}(H) \cup \{\{t\}, \{t^{-1}\}\}$ be the set of labels of the transitions of the fta \mathcal{B} . We can substitute in \mathcal{B} every L -labelled edge with $L \in \text{Rat}(H)$ by a finite automaton over the generating set X of H . In this way, we see that

$$L(\mathcal{B}) \in \text{Rat}(H * \{t, t^{-1}\}^*). \quad (66)$$

As every t -sequence is equivalent (modulo \approx) with at least one reduced sequence, we have

$$K = \pi_G(\pi_G^{-1}(K) \cap \text{Red}(H, t)). \quad (67)$$

With (65) we get

$$K = \pi_G(L(\mathcal{B}) \cap \text{Red}(H, t)). \quad (68)$$

Since \mathcal{B} is \approx -compatible, we have

$$\pi_G(\mathbf{L}(\mathcal{B})) \subseteq \pi_G([\mathbf{L}(\mathcal{B}) \cap \text{Red}(H, t)]_{\approx}) = \pi_G(\mathbf{L}(\mathcal{B}) \cap \text{Red}(H, t)).$$

Since also $\pi_G(\mathbf{L}(\mathcal{B}) \cap \text{Red}(H, t)) \subseteq \pi_G(\mathbf{L}(\mathcal{B}))$, (68) implies

$$K = \pi_G(\mathbf{L}(\mathcal{B})). \quad (69)$$

As any homomorphic image of a rational subset is rational, K is a rational subset of G . \square

4.5. Transfer results for rational sets

In this section we prove the following theorem, which allows to transfer important properties from $\text{Rat}(H)$ to $\text{Rat}(G)$.

Theorem 37. *Let $G = \langle H, t; t^{-1}at = \varphi(a)(a \in A) \rangle$, where H is a countably generated monoid with finite subgroups A and B , and $\varphi : A \rightarrow B$ is an isomorphism. Then we have:*

- (1) *If $\text{Rat}(H)$ is closed under intersection, then so is $\text{Rat}(G)$.*
- (2) *If $\text{Rat}(H)$ is closed under complement, then so is $\text{Rat}(G)$.*
- (3) *If the membership problem for rational subsets of H is decidable, then so is it for G .*
- (4) *If the emptiness problem for boolean combinations of rational subsets of H is decidable, then so is it for G .*

Proof. For statement (1), let $K_1, K_2 \in \text{Rat}(G)$ and $K = K_1 \cap K_2$. By Proposition 33 there are partitioned, \approx -compatible, and \sim -saturated fta \mathcal{A}_1 and \mathcal{A}_2 with labelling set $\text{Rat}(H)$ such that

$$\pi_G^{-1}(K_i) \cap \text{Red}(H, t) = \mathbf{L}(\mathcal{A}_i) \cap \text{Red}(H, t)$$

for $i \in \{1, 2\}$. Hence,

$$\begin{aligned} \pi_G^{-1}(K) \cap \text{Red}(H, t) &= \pi_G^{-1}(K_1) \cap \pi_G^{-1}(K_2) \cap \text{Red}(H, t) \\ &= \mathbf{L}(\mathcal{A}_1) \cap \mathbf{L}(\mathcal{A}_2) \cap \text{Red}(H, t). \end{aligned}$$

Let us consider the partitioned fta

$$\mathcal{A} = \mathcal{A}_1 \times \mathcal{A}_2.$$

By Lemma 15, $\mathbf{L}(\mathcal{A}) = \mathbf{L}(\mathcal{A}_1) \cap \mathbf{L}(\mathcal{A}_2)$, and by Lemma 17, \mathcal{A} is \approx -compatible and \sim -saturated. Moreover, since $\text{Rat}(H)$ is closed under intersection, the labelling set of \mathcal{A} is still $\text{Rat}(H)$. Hence K fulfils point (3) of Proposition 33. By point (1) of Proposition 33, K is a rational subset of G .

For statement (2), let $K \in \text{Rat}(G)$. By point (3) of Proposition 33 we have

$$\pi_G^{-1}(K) \cap \text{Red}(H, t) = \mathbf{L}(\mathcal{B}) \cap \text{Red}(H, t)$$

for a partitioned, \approx -compatible, and \sim -saturated fta \mathcal{B} with labelling set $\text{Rat}(H)$. By applying Proposition 28 to \mathcal{B} , we obtain

$$\pi_G^{-1}(K) \cap \text{Red}(H, t) = \text{L}(\mathcal{C})$$

for some partitioned, \sim -saturated, deterministic, and complete fta \mathcal{C} . The labelling set of this fta is $\text{Bool}(\text{Rat}(H))$, which equals $\text{Rat}(H)$, since $\text{Rat}(H)$ is supposed to be closed under complement.

Now take the complement automaton $\overline{\mathcal{C}}$ of \mathcal{C} , see Definition 19. By Lemma 20 it is a partitioned and \sim -saturated fta with

$$\text{L}(\overline{\mathcal{C}}) = \text{Red}(H, t) \setminus \text{L}(\mathcal{C}) \subseteq \text{Red}(H, t).$$

In particular, $\overline{\mathcal{C}}$ is also \approx -compatible. We obtain

$$\begin{aligned} \pi_G^{-1}(G \setminus K) \cap \text{Red}(H, t) &= (H * \{t, t^{-1}\}^* \setminus \pi_G^{-1}(K)) \cap \text{Red}(H, t) \\ &= \text{Red}(H, t) \setminus (\pi_G^{-1}(K) \cap \text{Red}(H, t)) \\ &= \text{Red}(H, t) \setminus \text{L}(\mathcal{C}) \\ &= \text{L}(\overline{\mathcal{C}}) \\ &= \text{L}(\overline{\mathcal{C}}) \cap \text{Red}(H, t). \end{aligned}$$

Hence, Proposition 33 implies that $G \setminus K$ is a rational subset of G . This proves statement (2).

For statement (3) suppose that the membership problem for rational subsets of H is decidable. Let $K = \pi(\text{L}(\mathcal{A}))$ for a given finite automaton \mathcal{A} over the free monoid $(X \cup \{t, t^{-1}\})^*$. By point (3) of Proposition 33 we have

$$\pi_G^{-1}(K) \cap \text{Red}(H, t) = \text{L}(\mathcal{B}) \cap \text{Red}(H, t)$$

for a partitioned, \approx -compatible, and \sim -saturated fta \mathcal{B} with labelling set $\text{Rat}(H)$. Moreover, since the membership problem for languages in $\text{Rat}(H)$ is decidable, \mathcal{B} can be constructed effectively from \mathcal{A} .

Now, to a given word $u \in (X \cup \{t, t^{-1}\})^*$ we first apply (effectively) as long as possible the rules of the semi-Thue system ST in order to obtain a word $v \in (X \cup \{t, t^{-1}\})^*$ of the form

$$v = v_0 t^{\alpha_1} v_1 \cdots t^{\alpha_i} v_i \cdots t^{\alpha_p} v_p$$

where $p \geq 0$, $\alpha_i \in \{1, -1\}$, $v_i \in X^*$, and $\pi_t(v) \in \text{Red}(H, t)$. Note that ST contains infinitely many rules, but nevertheless we can compute effectively such a word v : since the word problem is decidable for H , we can decide whether $[w]_{\equiv_H} \in A$ (resp. $[w]_{\equiv_H} \in B$ for a given word $w \in X^*$; this allows us to apply rules from ST in an effective way.

Since the membership problem is decidable for the labelling sets of \mathcal{B} , we can compute all the binary relations $\mu_{\mathcal{B}}(\pi_t(v_i))$, $\mu_{\mathcal{B}}(t)$, and $\mu_{\mathcal{B}}(t^{-1})$. This allows us to compute the binary relation $\mu_{\mathcal{B}}(\pi_t(v))$. Since $u \equiv v$, we have $\pi(u) \in K$ if and only if $\pi(v) \in K$ if and only if $\pi_t(v) \in \pi_G^{-1}(K)$ if and only if $\pi_t(v) \in \text{L}(\mathcal{B})$ if and only if $\mu_{\mathcal{B}}(\pi_t(v)) \cap \mathbf{1}_{\mathcal{B}} \times \mathbf{T}_{\mathcal{B}} \neq \emptyset$.

Let us finally prove statement (4). Assume that the emptiness problem for boolean combinations of rational subsets of H is decidable. Let K_1, \dots, K_p be rational subsets of G defined as $K_i = \pi(L(\mathcal{A}_i))$ for some given finite automaton \mathcal{A}_i over the free monoid $(X \cup \{t, t^{-1}\})^*$. Following the construction described for statement (2) of Theorem 37, we can construct effectively for every $i \in \{1, \dots, p\}$ a partitioned, \sim -saturated, deterministic, and complete fta \mathcal{C}_i , whose labelling set is $\text{Bool}(\text{Rat}(H))$, and such that

$$L(\mathcal{C}_i) = \pi_G^{-1}(K_i) \cap \text{Red}(H, t).$$

The boolean operations \cap and complement over \sim -saturated subsets of $\text{Red}(H, t)$ can be translated into the product and complement operation, respectively, over partitioned, \sim -saturated, deterministic, and complete fta reading only reduced sequences (for products, we have to use Lemma 17 and 18, for complements we use Lemma 20). Hence, from a given boolean combination K of the sets K_i , one can compute effectively a corresponding fta \mathcal{C} , whose labelling set is $\text{Bool}(\text{Rat}(H))$, and such that

$$L(\mathcal{C}) = \pi_G^{-1}(K) \cap \text{Red}(H, t).$$

Emptiness of the set K reduces to emptiness of $L(\mathcal{C})$, which itself reduces to the emptiness problem for the label sets of \mathcal{C} . Hence, under the hypothesis that emptiness is decidable for languages in $\text{Bool}(\text{Rat}(H))$, we can test whether $L(\mathcal{C}) = \emptyset$ or not. \square

Remark 38. For a given monoid \mathbb{M} , closure of $\text{Rat}(\mathbb{M})$ under complement implies closure of $\text{Rat}(\mathbb{M})$ under intersection, but the converse does not hold in general: we give examples of such monoids \mathbb{M} where $\text{Rat}(\mathbb{M})$ is closed under intersection but not under complement in Appendix B. This shows that points (1) and (2) of Theorem 37 cannot be deduced in a straightforward manner one from the other.

The same is also true for (3) and (4): For the group $F \times \mathbb{Z}$ (where F is the free group of rank 2), the membership problem for rational subsets is decidable [KSS06], whereas the inclusion problem for rational subsets (which is a special instance of the emptiness problem for boolean combinations of rational subsets) is undecidable [AH89].

4.6. Transfer results for finitely generated subgroups

We suppose in this section that $H = X^*/\equiv$ (and hence G) is a group. We assume moreover that the generating set X is endowed with an involution $x \mapsto x^{-1}$ such that $xx^{-1} \equiv x^{-1}x \equiv \varepsilon$. The following subset of $\mathcal{P}(H)$ will serve for labelling fta in the sequel:

$$\mathcal{S} = \{Kh \mid K \text{ is a finitely generated subgroup of } H, h \in H\}$$

Note that every singleton subset $\{h\}$ belongs to \mathcal{S} .

4.6.1. Normalization of automata

Let us prove first a normalization lemma for finitely generated subgroups of G , which parallels Proposition 33:

Proposition 39. *Let K be a subgroup of G . The following are equivalent:*

- (1) K is finitely generated.
- (2) K is rational.
- (3) There exists some finite automaton \mathcal{A} over the free monoid $(X \cup \{t, t^{-1}\})^*$ such that $K = \pi(\mathbf{L}(\mathcal{A}))$.
- (4) $\pi_G^{-1}(K) \cap \text{Red}(H, t) = \mathbf{L}(\mathcal{B}) \cap \text{Red}(H, t)$ for some partitioned, \approx -compatible, and \sim -saturated fta \mathcal{B} with labelling set \mathcal{S} .

Moreover, in (4), if the generalized word problem of H is decidable, then \mathcal{B} can be constructed effectively from \mathcal{A} in (3).

Proof. We follow the arguments from the proof of Proposition 33.

(1) \Leftrightarrow (2): This is just Theorem 1.

(2) \Leftrightarrow (3): This has been shown in Proposition 33.

(3) \Rightarrow (4): We must here slightly adapt the corresponding proof given for Proposition 33. Let us start with a finite automaton $\mathcal{A} = \langle \mathcal{X}, \mathbf{Q}, \delta, \mathbf{l}, \mathbf{T} \rangle$ over the free monoid $(X \cup \{t, t^{-1}\})^*$ such that

$$K = \pi(\mathbf{L}(\mathcal{A})).$$

Claim 40. *There exists a finite automaton \mathcal{A}_1 over the free monoid $(X \cup \{t, t^{-1}\})^*$ such that*

$$\mathbf{L}(\mathcal{A}) \subseteq \mathbf{L}(\mathcal{A}_1) = \mathbf{L}(\mathcal{A}_1) \rightarrow_{\mathcal{S}\mathbf{T}}^* \subseteq [\mathbf{L}(\mathcal{A})]_{\equiv}. \quad (70)$$

Moreover, if the generalized word problem of H is decidable, then \mathcal{A}_1 can be constructed effectively from \mathcal{A} .

We assume that \mathcal{A} is trim, i.e., every state of \mathcal{A} is accessible from an initial state and co-accessible from a final state. We start with

$$\delta_1 = \delta \cup \{(r, c^{-1}, q) \mid c \in X \cup \{t, t^{-1}\}, (q, c, r) \in \delta\}.$$

Since $\pi(\mathbf{L}(\mathcal{A}))$ is a group, we have

$$\mathbf{L}(\mathcal{A}) \subseteq \mathbf{L}(\langle \mathcal{X}, \mathbf{Q}, \delta_1, \mathbf{l}, \mathbf{T} \rangle) \subseteq [\mathbf{L}(\mathcal{A})]_{\equiv}.$$

For all states $q, q' \in \mathbf{Q}$ we define $\ell(\delta_1, q, q')$ as a shortest word over X which labels a δ_1 -path from q to q' . We saturate δ_1 under the following inference rules, where

$u \in X^*$:

$$\frac{p \xrightarrow{t^{-1}}_{\delta_1} q \xrightarrow{u}_{\delta_1} q \xrightarrow{\ell(\delta_1, q, q')}_{\delta_1} q' \xrightarrow{t}_{\delta_1} r \wedge u\ell(\delta_1, q, q') \equiv_H a \in A}{p \xrightarrow{\varphi(a)}_{\delta_1} r} \quad (71)$$

$$\frac{p \xrightarrow{t}_{\delta_1} q \xrightarrow{u}_{\delta_1} q \xrightarrow{\ell(\delta_1, q, q')}_{\delta_1} q' \xrightarrow{t^{-1}}_{\delta_1} r \wedge u\ell(\delta_1, q, q') \equiv_H b \in B}{p \xrightarrow{\varphi^{-1}(b)}_{\delta_1} r} \quad (72)$$

Let δ_1 be the resulting transition relation and let

$$\mathcal{A}_1 = \langle \mathcal{X} \cup A \cup B, \mathbf{Q}, \delta_1, \mathbf{l}, \mathbf{T} \rangle.$$

It satisfies (70). For this notice that if $q \xrightarrow{v}_{\delta_1} q'$ for a word $v \in X^*$ with $v \equiv_H a \in A$ then $v\ell(\delta_1, q, q')^{-1}\ell(\delta_1, p, q) \equiv_H a$ and $q \xrightarrow{v\ell(\delta_1, q, q')^{-1}}_{\delta_1} q$, i.e., the word $v\ell(\delta_1, q, q')^{-1}$ is a candidate for u in the above rule (71).

It remains to show the effectiveness statement in Claim 40. Let us consider for every value of δ_1 and every $q \in \mathbf{Q}$ the set

$$P_1(\delta_1, q) = \{u \in X^* \mid q \xrightarrow{u}_{\delta_1} q\}.$$

By Theorem 1, $\pi_t(P_1(\delta_1, q))$ is a finitely generated subgroup of H , and by the proof given in Section 2.2, a finite set of generators for it can be explicitly computed. If the generalized word problem of H is decidable, then the validity of the hypothesis of the inference rules (71) and (72) can be algorithmically tested so that δ_1 can be algorithmically computed. This proves Claim 40.

Claim 41. *There exists a \approx -compatible fta \mathcal{A}_2 over $H * \{t, t^{-1}\}^*$ with labelling set \mathcal{S} such that*

$$L(\mathcal{A}_2) = \pi_t(L(\mathcal{A}_1)). \quad (73)$$

Moreover, \mathcal{A}_2 can be constructed effectively from \mathcal{A}_1 .

Analogously to the proof of Claim 35, we do a collecting process:

$$\begin{aligned} \delta_2 = & \{(q, t^\alpha, r) \mid \alpha \in \{1, -1\}, (q, t^\alpha, r) \in \delta_1\} \cup \\ & \{(q, \pi_t(P_1(\delta_1, q)\ell(\delta_1, q, r)), r) \mid q, r \in \mathbf{Q}\}. \end{aligned} \quad (74)$$

This step can be shown to be effective by the arguments already used for the proof of Claim 40. Clearly (73) holds. Moreover, (70) implies that $L(\mathcal{A}_1) \subseteq [L(\mathcal{A}_1) \cap \text{Irr}(\text{ST})]_{\equiv}$. Now $L(\mathcal{A}_2) \subseteq [L(\mathcal{A}_2) \cap \text{Red}(H, t)]_{\approx}$ can be shown in the same way as in the proof of Claim 35. Hence, \mathcal{A}_2 is \approx -compatible.

Claim 42. *There exists a partitioned and \approx -compatible fta \mathcal{A}_3 over $H * \{t, t^{-1}\}^*$ with labelling set \mathcal{S} and such that*

$$L(\mathcal{A}_3) = L(\mathcal{A}_2).$$

Moreover, \mathcal{A}_3 can be constructed effectively from \mathcal{A}_2 .

As for Claim 36 set $\mathcal{A}_3 = \mathcal{A}_2 \times \mathcal{G}_6$.

Let us now apply Proposition 22 to the set $\mathcal{F} = \mathcal{S}$ and the fta \mathcal{A}_3 , and let \mathcal{B} be the resulting partitioned, \approx -compatible, and \sim -saturated fta. Note that the labelling set of \mathcal{B} is still \mathcal{S} : If Kh labels a transition of \mathcal{A}_3 , where K is a finitely generated subgroup of H and $h \in H$, then a transition of \mathcal{B} may be labelled with $c(Kh)c'$, where $c, c' \in A \cup B$. But we have $c(Kh)c' = (cKc^{-1})(chc') \in \mathcal{S}$.

We obtain the following relationships between the automata \mathcal{A} , \mathcal{A}_1 , \mathcal{A}_2 , \mathcal{A}_3 , and \mathcal{B} :

$$K = \pi(\mathbb{L}(\mathcal{A})) \subseteq G \quad (75)$$

$$\mathbb{L}(\mathcal{A}) \subseteq \mathbb{L}(\mathcal{A}_1) \subseteq [\mathbb{L}(\mathcal{A})]_{\equiv} \subseteq (X \cup \{t, t^{-1}\})^* \quad (76)$$

$$\mathbb{L}(\mathcal{A}_2) = \pi_t(\mathbb{L}(\mathcal{A}_1)) \subseteq H * \{t, t^{-1}\}^* \quad (77)$$

$$\mathbb{L}(\mathcal{A}_3) = \mathbb{L}(\mathcal{A}_2) \subseteq H * \{t, t^{-1}\}^* \quad (78)$$

$$\mathbb{L}(\mathcal{B}) \cap \text{Red}(H, t) = [\mathbb{L}(\mathcal{A}_3)]_{\approx} \cap \text{Red}(H, t) \subseteq H * \{t, t^{-1}\}^* \quad (79)$$

From (76) we obtain $[\mathbb{L}(\mathcal{A})]_{\equiv} = [\mathbb{L}(\mathcal{A}_1)]_{\equiv}$. Now, we can deduce $\pi_G^{-1}(K) \cap \text{Red}(H, t) = \mathbb{L}(\mathcal{B}) \cap \text{Red}(H, t)$ in the same way as in the corresponding part of the proof of Proposition 33.

(4) \Rightarrow (2): This follows from Proposition 33 since, $\mathcal{S} \subseteq \text{Rat}(H)$. \square

Using Proposition 39, we can next prove a transfer theorem for finitely generated subgroups:

Theorem 43. *Let $G = \langle H, t; t^{-1}at = \varphi(a)(a \in A) \rangle$, where H is a countably generated group with finite subgroups A and B , and $\varphi : A \rightarrow B$ is an isomorphism. Then we have:*

- (1) *If H has a decidable generalized word problem, then so has G .*
- (2) *If H has the Howson property, then so has G .*
- (3) *If H has the LERF property, then so has G .*

Proof. For property (1), let $u_1, \dots, u_n, u \in (X \cup \{t, t^{-1}\})^*$. Let $K \subseteq G$ be the subgroup generated by $\{\pi(u_1), \dots, \pi(u_n)\}$. We give an algorithm for deciding whether $\pi(u) \in K$ or not:

Step 1: We construct from u_1, \dots, u_n a finite automaton \mathcal{A} over the free monoid $(X \cup \{t, t^{-1}\})^*$ such that

$$K = \pi(\mathbb{L}(\mathcal{A})).$$

Step 2: Using Proposition 39 we can compute from \mathcal{A} effectively a partitioned, \approx -compatible, and \sim -saturated fta \mathcal{B} with labelling set \mathcal{S} such that

$$\pi_G^{-1}(K) \cap \text{Red}(H, t) = \mathbb{L}(\mathcal{B}) \cap \text{Red}(H, t).$$

Step 3: Starting from the word $u \in (X \cup \{t, t^{-1}\})^*$ and applying (effectively) the rules of ST as long as possible, one computes a word $v \in (X \cup \{t, t^{-1}\})^*$ of the form

$$v = v_0 t^{\alpha_1} v_1 \cdots t^{\alpha_i} v_i \cdots t^{\alpha_p} v_p$$

where $p \geq 0$, $\alpha_i \in \{1, -1\}$, $v_i \in X^*$, and $\pi_t(v) \in \text{Red}(H, t)$.

Step 4: From \mathcal{B} one can compute all the binary relations $\mu_{\mathcal{B}}(\pi_t(v_i))$, $\mu_{\mathcal{B}}(t)$, and $\mu_{\mathcal{B}}(t^{-1})$. This allows to compute the binary relation $\mu_{\mathcal{B}}(\pi_t(v))$. Since $u \equiv v$, we have $\pi(u) \in K$ if and only if $\pi(v) \in K$ if and only if $\pi_t(v) \in \pi_G^{-1}(K)$ if and only if $\pi_t(v) \in L(\mathcal{B})$ if and only if $\mu_{\mathcal{B}}(\pi_t(v)) \cap \mathbb{1}_{\mathcal{B}} \times \mathbb{T}_{\mathcal{B}} \neq \emptyset$.

Let us now prove (2): Suppose that H has the Howson property. Let K_1 and K_2 be two finitely generated subgroups of G . By Proposition 39, there exist partitioned, \approx -compatible, and \sim -saturated fta \mathcal{B}_1 and \mathcal{B}_2 with labelling set \mathcal{S} such that

$$\pi_G^{-1}(K_i) \cap \text{Red}(H, t) = L(\mathcal{B}_i) \cap \text{Red}(H, t)$$

for $i \in \{1, 2\}$. Let

$$\mathcal{B} = \mathcal{B}_1 \times \mathcal{B}_2.$$

By the same arguments as for point (1) of Theorem 37, \mathcal{B} is partitioned, \approx -compatible, and \sim -saturated. Let us check that its transition labels belong to \mathcal{S} : Every label L of \mathcal{B} either belongs to $\{\{t\}, \{t^{-1}\}\}$ or has the form

$$L = F_1 h_1 \cap F_2 h_2 \neq \emptyset$$

for finitely generated subgroups $F_1, F_2 \leq H$ and $h_1, h_2 \in H$. Since $F_1 h_1 \cap F_2 h_2 \neq \emptyset$, there exist $f_1 \in F_1$ and $f_2 \in F_2$ such that $f_1 h_1 = f_2 h_2$ in H . We thus have

$$\begin{aligned} F_1 h_1 \cap F_2 h_2 &= ((F_1 f_1^{-1} f_2) \cap F_2) h_2 \\ &= (F_1 f_2 \cap F_2) h_2 \\ &= (F_1 \cap F_2) f_2 h_2. \end{aligned}$$

By the Howson property for H , $F_1 \cap F_2$ is a finitely generated subgroup of H . Hence, $(F_1 \cap F_2) f_2 h_2$ belongs to \mathcal{S} . Thus

$$\pi_G^{-1}(K_1 \cap K_2) \cap \text{Red}(H, t) = L(\mathcal{B}) \cap \text{Red}(H, t)$$

for a partitioned, \approx -compatible, and \sim -saturated fta \mathcal{B} with labelling set \mathcal{S} . By Proposition 39 this implies that $K_1 \cap K_2$ is finitely generated.

Finally, we prove point (3). In order to treat the LERF property (see Definition 3) we have to introduce a technical notion about finite t -automata, which is stronger than \approx -compatibility:

Definition 44. Let $\mathcal{B} = \langle \mathcal{L}, \mathcal{Q}, \tau, \delta, \mathbb{1}, \mathbb{T} \rangle$ be a partitioned fta. It is said to be downwards-closed (briefly d -closed) if for all $q_1, q_2, q_3, q_4, q_5, q_6 \in \mathcal{Q}$, $h_2, h_6 \in H$, $a \in A$, and $b \in B$ such that $\tau(q_1) \neq \tau(q_2)$ and $\tau(q_5) \neq \tau(q_6)$:

$$q_1 \xrightarrow{h_2}_{\mathcal{B}} q_2 \xrightarrow{t^{-1}}_{\mathcal{B}} q_3 \xrightarrow{a}_{\mathcal{B}} q_4 \xrightarrow{t}_{\mathcal{B}} q_5 \xrightarrow{h_6}_{\mathcal{B}} q_6 \Rightarrow q_1 \xrightarrow{h_2 \varphi(a) h_6}_{\mathcal{B}} q_6 \quad (80)$$

$$q_1 \xrightarrow{h_2}_{\mathcal{B}} q_2 \xrightarrow{t}_{\mathcal{B}} q_3 \xrightarrow{b}_{\mathcal{B}} q_4 \xrightarrow{t^{-1}}_{\mathcal{B}} q_5 \xrightarrow{h_6}_{\mathcal{B}} q_6 \Rightarrow q_1 \xrightarrow{h_2 \varphi^{-1}(b) h_6}_{\mathcal{B}} q_6 \quad (81)$$

Lemma 45. *A partitioned and d -closed fta is \approx -compatible.*

Proof. Assume that the partitioned fta $\mathcal{B} = \langle \mathcal{L}, \mathcal{Q}, \tau, \delta, \mathbf{l}, \mathbf{T} \rangle$ is d -closed and let $s \in \mathbf{L}(\mathcal{B})$. We have to show that $s \in [\mathbf{L}(\mathcal{B}) \cap \text{Red}(H, t)]_{\approx}$. For this, assume that s is not reduced, i.e., $s = s_1 t^{-1} a t s_2$. Since $s \in \mathbf{L}(\mathcal{B})$, there exists a path in \mathcal{B} , labelled with s , leading from a state of type $(1, H)$ to a state of type $(1, 1)$. Since \mathcal{B} is partitioned, the sequences s_1 and s_2 cannot be empty: note that no t -labelled (resp. t^{-1} -labelled) transitions exits (resp. enters) a state of type $(1, H)$ (resp. $(1, 1)$) in a partitioned fta. Hence we have $s = s'_1 h_2 t^{-1} a t h_6 s'_2$ for some $h_2, h_6 \in H$. Moreover, there exist states $q_1, q_2, q_3, q_4, q_5, q_6 \in \mathcal{Q}$ such that (80) holds. Since \mathcal{B} is partitioned and the path labelled with s leads from a state of type $(1, H)$ to a state of type $(1, 1)$, we have $\tau(q_1) \neq \tau(q_2)$ and $\tau(q_5) \neq \tau(q_6)$ (see the last statement in Lemma 7). Hence, since \mathcal{B} is d -closed, we have $q_1 \xrightarrow{h_2 \varphi(a) h_6}_{\mathcal{B}} q_6$, i.e., $s'_1 (h_2 \varphi(a) h_6) s'_2 \in \mathbf{L}(\mathcal{B})$. By continuing in this way, we finally arrive at a reduced sequence $s' \in \mathbf{L}(\mathcal{B}) \cap \text{Red}(H, t)$ such that $s \approx s'$. \square

Let us now continue with the proof for point (3) in Theorem 43. Assume that H has the LERF property. We will use the alternative definition from Remark 4. Let $g \in G$, let $K \leq G$ be some finitely generated subgroup of G , and let us suppose that

$$g \notin K. \quad (82)$$

By Proposition 39, there exists a partitioned, \approx -compatible, and \sim -saturated fta

$$\mathcal{B} = \langle \mathcal{L}_{\mathcal{B}}, \mathcal{Q}_{\mathcal{B}}, \tau_{\mathcal{B}}, \delta_{\mathcal{B}}, \mathbf{l}_{\mathcal{B}}, \mathbf{T}_{\mathcal{B}} \rangle$$

with labelling set \mathcal{S} such that

$$\pi_G^{-1}(K) \cap \text{Red}(H, t) = \mathbf{L}(\mathcal{B}) \cap \text{Red}(H, t). \quad (83)$$

Claim 46. *The fta \mathcal{B} can be chosen to be d -closed.*

Let us look at the fta \mathcal{A}_2 produced in the proof of Claim 41 by equation (74). This automaton is not partitioned but has both properties (80) and (81). The partitioned automaton $\mathcal{A}_3 = \mathcal{A}_2 \times \mathcal{G}_6$ still satisfies (80) and (81), since for all q_1, \dots, q_6 fulfilling the hypothesis of (80) and (81), respectively, the edge $(\tau_{\mathcal{A}_2}(q_1), H, \tau_{\mathcal{A}_2}(q_6))$ belongs to \mathcal{E}_6 . Let us look finally at the fta \mathcal{B} obtained from \mathcal{A}_3 by the construction in the proof of Proposition 22. Suppose that the hypothesis of implication (80) is fulfilled, i.e.,

$$q_1 g_1 \xrightarrow{h_2}_{\mathcal{B}} q_2 g_2 \xrightarrow{t^{-1}}_{\mathcal{B}} q_3 g_3 \xrightarrow{a}_{\mathcal{B}} q_4 g_4 \xrightarrow{t}_{\mathcal{B}} q_5 g_5 \xrightarrow{h_6}_{\mathcal{B}} q_6 g_6 \quad (84)$$

where $q_i \in \mathcal{Q}_{\mathcal{A}_3}$, $g_i \in \gamma(q_i)$, $\tau_{\mathcal{B}}(q_1 g_1) \neq \tau_{\mathcal{B}}(q_2 g_2)$, $\tau_{\mathcal{B}}(q_5 g_5) \neq \tau_{\mathcal{B}}(q_6 g_6)$, and hence $\tau_{\mathcal{A}_3}(q_1) \neq \tau_{\mathcal{A}_3}(q_2)$, $\tau_{\mathcal{A}_3}(q_5) \neq \tau_{\mathcal{A}_3}(q_6)$. Assembling five of the diagrams described in Figure 4, we obtain from (84) in the fta \mathcal{A}_3 the path

$$q_1 \xrightarrow{g_1 h_2 g_2^{-1}}_{\mathcal{A}_3} q_2 \xrightarrow{t^{-1}}_{\mathcal{A}_3} q_3 \xrightarrow{g_3 a g_4^{-1}}_{\mathcal{A}_3} q_4 \xrightarrow{t}_{\mathcal{A}_3} q_5 \xrightarrow{g_5 h_6 g_6^{-1}}_{\mathcal{A}_3} q_6,$$




Fig. 5. \mathcal{B} is d -closed

where

$$\varphi^{-1}(g_2) = g_3 \quad \text{and} \quad \varphi(g_4) = g_5. \quad (86)$$

Since \mathcal{A}_3 is d -closed, we get

$$q_1 \xrightarrow{(g_1h_2g_2^{-1})\varphi(g_3ag_4^{-1})(g_5h_6g_6^{-1})} \mathcal{A}_3 q_6,$$

i.e., with (86),

$$q_1 \xrightarrow{(g_1h_2)\varphi(a)(h_6g_6^{-1})} \mathcal{A}_3 q_6.$$

Using the left diagram in Figure 4, we obtain

$$q_1g_1 \xrightarrow{h_2\varphi(a)h_6} \mathcal{B} q_6g_6.$$

Thus, implication (80) holds, see also Figure 5. A similar verification can be done for implication (81). Claim 46 is thus established.

Let

$$\mathcal{L}_{\mathcal{B}} \cap \mathcal{P}(H) = \{F_0r_0, \dots, F_\ell r_\ell\}$$

be the set of H -labels of \mathcal{B} , where $F_i \leq H$ is finitely generated and $r_i \in H$. Let

$$s = h_0t^{\alpha_1}h_1 \cdots t^{\alpha_n}h_n \in \text{Red}(H, t)$$

be a reduced sequence such that

$$\pi_G(s) = g \notin K. \quad (87)$$

By (83), (87), and the fact that s is reduced, we have

$$s \notin \mathbf{L}(\mathcal{B}). \quad (88)$$

For every subgroup $N' \leq H$ let us consider the following properties:

$$N' \cap A = 1 = N' \cap B \quad (89)$$

$$\forall i \in \{0, \dots, \ell\} \forall c \in A \cup B : c \in N' F_i r_i \Leftrightarrow c \in F_i r_i \quad (90)$$

$$\forall i \in \{0, \dots, \ell\}, j \in \{0, \dots, n\} : h_j \in N' F_i r_i \Leftrightarrow h_j \in F_i r_i \quad (91)$$

$$\forall j \in \{0, \dots, n\} : h_j \in N' A \Leftrightarrow h_j \in A \wedge h_j \in N' B \Leftrightarrow h_j \in B \quad (92)$$

Notice that the directions \Leftarrow in (90)–(92) is trivially satisfied for every subgroup N' . As H meets the LERF property, for each value of $i \in \{0, \dots, \ell\}, j \in \{0, \dots, n\}$, and each property P among the above properties (89)–(92), there exists some normal subgroup of finite index $N(i, j, P) \trianglelefteq H$ fulfilling the given property. For instance, for (90) one can argue as follows: if $c \notin F_i r_i$, then $cr_i^{-1} \notin F_i$. Since F_i is finitely generated, the LERF property for H implies the existence of a normal subgroup $N' \leq H$ of finite index in H such that $cr_i^{-1} \notin N' F_i$, i.e., $c \notin N' F_i r_i$.

Let us define

$$N = \bigcap_{i,j,P} N(i, j, P).$$

This subgroup N is normal and has finite index in H . Moreover, since every property among (89)–(92) is *hereditary* in the sense that if $N_1 \leq N_2 \leq H$ and N_2 meets the property, then N_1 meets the same property as well, N fulfils simultaneously all the properties (89)–(92). Let us set

$$G' = \langle H, t; x = 1 (x \in N), t^{-1}at = \varphi(a) (a \in A) \rangle.$$

Let $\pi' : H \rightarrow H' = H/N$ be the canonical projection. Due to (89), this morphism π' fulfils condition (19) from Section 3.1. Thus, the group G' is also the HNN-extension of $H' = H/N$ with stable letter t , associated subgroups A, B , and the partial isomorphism φ :

$$G' = \langle H/N, t; t^{-1}at = \varphi(a) (a \in A) \rangle.$$

Let us use the notation from Section 3.1. Let $\pi'_*(\mathcal{B})$ be the \sim' -saturated fta from Lemma 21, which satisfies

$$L(\pi'_*(\mathcal{B})) = \pi'_*(L(\mathcal{B})).$$

Let us define:

$$s' = \pi'_*(s) = \pi'(h_0)t^{\alpha_1}\pi'(h_1) \cdots t^{\alpha_n}\pi'(h_n). \quad (93)$$

Since s is a reduced sequence in $H * \{t, t^{-1}\}^*$, condition (92) implies

$$s' \in \text{Red}(H', t). \quad (94)$$

By (88) and (91) we have

$$s' \notin L(\pi'_*(\mathcal{B})). \quad (95)$$

By (83) and the fact that \mathcal{B} is \approx -compatible, we get

$$\begin{aligned}
 \pi_G(\mathbf{L}(\mathcal{B})) &= \pi_G([\mathbf{L}(\mathcal{B})]_{\approx}) \\
 &= \pi_G([\mathbf{L}(\mathcal{B}) \cap \text{Red}(H, t)]_{\approx}) \\
 &= \pi_G(\mathbf{L}(\mathcal{B}) \cap \text{Red}(H, t)) \\
 &= \pi_G(\pi_G^{-1}(K) \cap \text{Red}(H, t)) \\
 &= K,
 \end{aligned}$$

and thus,

$$\pi_{G'}^{-1}(\pi'(K)) = \pi_{G'}^{-1}(\pi'(\pi_G(\mathbf{L}(\mathcal{B}))).$$

The commutative diagram in Figure 1 yields

$$\pi_{G'}^{-1}(\pi'(K)) = \pi_{G'}^{-1}(\pi_{G'}(\pi'_*(\mathbf{L}(\mathcal{B})))) = [\pi'_*(\mathbf{L}(\mathcal{B}))]_{\approx'} = [\mathbf{L}(\pi'_*(\mathcal{B}))]_{\approx'}.$$

By Claim 46 and (90), $\pi'_*(\mathcal{B})$ is d -closed and hence \approx' -compatible by Lemma 45. Hence,

$$\begin{aligned}
 \pi_{G'}^{-1}(\pi'(K)) \cap \text{Red}(H', t) &= [\mathbf{L}(\pi'_*(\mathcal{B}))]_{\approx'} \cap \text{Red}(H', t) \\
 &= [\mathbf{L}(\pi'_*(\mathcal{B}) \cap \text{Red}(H', t))]_{\approx'} \cap \text{Red}(H', t) \\
 &= [\mathbf{L}(\pi'_*(\mathcal{B})) \cap \text{Red}(H', t)]_{\sim'} \\
 &= \mathbf{L}(\pi'_*(\mathcal{B})) \cap \text{Red}(H', t),
 \end{aligned}$$

where the last equality holds since $\pi'_*(\mathcal{B})$ is \sim' -saturated by Lemma 21. Together with (94) and (95) we get

$$s' \notin \pi_{G'}^{-1}(\pi'(K)).$$

This shows that $\pi_{G'}(s') \notin \pi'(K)$, i.e., by (93), $\pi_{G'}(\pi'_*(s)) \notin \pi'(K)$. The commutative diagram in Figure 1 implies $\pi'(\pi_G(s)) \notin \pi'(K)$, i.e., by (87),

$$\pi'(g) \notin \pi'(K). \tag{97}$$

The group G' is the fundamental group of a graph of groups (with just one vertex and one edge) with finite vertex groups (namely H/N), hence it is a virtually free group, i.e., it has a free subgroup of finite index, see e.g. [DD90, Corollary IV.1.9, p.104]. By the result of [Hal49] every free group is LERF and by [Sco78, Lemma 1.1, p. 556] every group with a finite index subgroup which is LERF is itself LERF. Hence G' is LERF. In particular, (97) implies the existence of a finite group G'' and a homomorphism $\pi'' : G' \rightarrow G''$ such that

$$\pi''(\pi'(g)) \notin \pi''(\pi'(K)). \tag{98}$$

By Remark 4, this proves that G is LERF.

5. Amalgamated free products

In the rest of this paper, we will prove for amalgamated free products results similar to those for HNN-extensions from Section 4.

Let us consider two monoids H_1 and H_2 , two finite *subgroups* $A_1 \leq H_1, A_2 \leq H_2$, and an isomorphism $\varphi : A_1 \rightarrow A_2$. The corresponding amalgamated free product

$$G = \langle H_1 * H_2; a = \varphi(a) (a \in A_1) \rangle \quad (99)$$

is defined by $G = (H_1 * H_2) / \approx$, where \approx is the congruence on the free product $H_1 * H_2$ generated by the equations $a = \varphi(a)$ for $a \in A_1$. An (H_1, H_2) -*sequence* is an element $s \in H_1 * H_2$; it has a unique decomposition of the form

$$s = h_0 k_1 h_1 \cdots k_i h_i \cdots k_n h_n, \quad (100)$$

where $n \geq 0, h_1, \dots, h_{n-1} \in H_2 \setminus 1, k_1, \dots, k_n \in H_1 \setminus 1$, and $h_0, h_n \in H_2$. A *reduced* (H_1, H_2) -*sequence* is an element $s \in H_1 * H_2$ of the form

$$s = h_0 k_1 h_1 \cdots k_i h_i \cdots k_n h_n, \quad (101)$$

where $n \geq 0, h_1, \dots, h_{n-1} \in H_2 \setminus A_2, k_1, \dots, k_n \in H_1 \setminus A_1$ and $h_0, h_n \in H_2$. We denote by $\text{Red}(H_1, H_2)$ the set of all reduced (H_1, H_2) -sequences.

Assume now that $H_1 = X_1^* / \equiv_{H_1}$ and $H_2 = X_2^* / \equiv_{H_2}$ for generating sets X_1 and X_2 such that $X_1 \cap X_2 = \emptyset$. Then $G = (X_1 \cup X_2)^* / \equiv$ where \equiv is the smallest congruence over the free monoid $(X_1 \cup X_2)^*$ such that $(\equiv_{H_1} \cup \equiv_{H_2}) \subseteq \equiv$ and $a \equiv \varphi(a)$ for all $a \in A_1$. Let

$$\pi : (X_1 \cup X_2)^* \rightarrow G$$

denote the canonical morphism defined by $\pi(w) = [w]_{\equiv}$ for $w \in (X_1 \cup X_2)^*$. This morphism naturally factorizes as

$$\pi = \pi_a \circ \pi_G$$

where

$$\begin{aligned} \pi_a : (X_1 \cup X_2)^* &\rightarrow H_1 * H_2 \text{ and} \\ \pi_G : H_1 * H_2 &\rightarrow G. \end{aligned}$$

The kernel of π_G coincides with the congruence \approx over $H_1 * H_2$ defined above.

We define the binary relation $\overset{1}{\sim}$ over $H_1 * H_2$ as follows: For all $s, s' \in H_1 * H_2$ let $s \overset{1}{\sim} s'$ if and only if there are (H_1, H_2) -sequences s_1 and s_2 and $a, b \in A_1$ such that either

$$\begin{aligned} \exists h \in H_1 : s = s_1 h s_2 \wedge s' = s_1 \varphi(a) (a^{-1} h b) \varphi(b^{-1}) s_2 \quad \text{or} \\ \exists h \in H_2 : s = s_1 h s_2 \wedge s' = s_1 a (\varphi(a^{-1}) h \varphi(b)) b^{-1} s_2. \end{aligned}$$


Let us denote by $\sim \subseteq \overset{1}{\sim}$ the congruence over $H_1 * H_2$ generated by $\overset{1}{\sim}$; it coincides with the reflexive and transitive closure of $\overset{1}{\sim}$. Equivalently, if

$$\begin{aligned} s &= h_0 k_1 h_1 k_2 \cdots h_{n-1} k_n h_n \text{ and} \\ s' &= h'_0 k'_1 h'_1 k'_2 \cdots h'_{m-1} k'_m h'_m \end{aligned}$$

(with $n, m \geq 0$, and $k_1, \dots, k_n, k'_1, \dots, k'_m \in H_1$, $h_0, \dots, h_n, h'_0, \dots, h'_m \in H_2$) are reduced sequences, then $s \sim s'$ if and only if $n = m$ and there exist $a_1, \dots, a_{2n} \in A_1$ such that:

- $h_i \varphi(a_{2i+1}) = \varphi(a_{2i}) h'_i$ in H_2 for $0 \leq i \leq n$ (here we set $a_0 = a_{2n+1} = 1$)
- $k_i a_{2i} = a_{2i-1} k'_i$ in H_1 for $1 \leq i \leq n$

In other words, there exists a Van Kampen diagram of the following kind, where $n = m = 4$. Here, we omit the $\varphi(a_i)$ -labelled edge, which runs parallel to the a_i -labelled vertical edge ($1 \leq i \leq 2n$).^c Light-shaded (resp. dark-shaded) areas represent relation in H_2 (resp. relations in H_1).



The set $\text{Red}(H_1, H_2)$ is again saturated by the congruence \sim , i.e., $s \sim s'$ implies $s \in \text{Red}(H_1, H_2) \Leftrightarrow s' \in \text{Red}(H_1, H_2)$. Here it is important that A_1 and A_2 are groups.

For the case that H_1 and H_2 (and hence G) are groups, it is well-known that G can be embedded into the HNN-extension

$$\widehat{G} = \langle H_1 * H_2, t; t^{-1}at = \varphi(a)(a \in A_1) \rangle$$

by the map Φ with

$$\Phi : k \mapsto t^{-1}kt \text{ for } k \in H_1 \text{ and } h \mapsto h \text{ for } h \in H_2, \quad (102)$$

see e.g. [LS77, Theorem 2.6, p. 187]. This result generalizes to our situation, where H_1 and H_2 are monoids but A_1 and A_2 are groups: Assume that $s_1, s_2 \in \text{Red}(H_1, H_2)$ such that $\Phi(s) \approx \Phi(s')$. Then $\Phi(s), \Phi(s') \in \text{Red}(H, t)$, and Lemma 5 implies $\Phi(s) \sim \Phi(s')$. The defining relations of \sim imply $s \sim s'$.

With our definitions of the relations \approx and \sim on the free product $H_1 * H_2$ and the new definition of a reduced sequence, Lemma 5 again holds:

Lemma 47. *Let $s, s' \in \text{Red}(H_1, H_2)$. Then $s \approx s'$ if and only if $s \sim s'$.*

Proof. Assume that $s \approx s'$ for $s, s' \in \text{Red}(H_1, H_2)$. The defining relations for \approx imply $\Phi(s) \approx \Phi(s')$. Then, the above arguments yield $s \sim s'$. \square

^cAlternatively, one may assume that $A_1 = A_2 = A$, $H_1 \cap H_2 = A$, where A is a subgroup of H_1 and H_2 , and that φ is the identity map. Then one obtains exactly the above Van Kampen diagram.

6. Finite automata for amalgamated free products

We fix throughout this section two monoids H_1, H_2 , two finite *subgroups* $A_1 \leq H_1, A_2 \leq H_2$, an isomorphism $\varphi : A_1 \rightarrow A_2$ and the corresponding amalgamated free product

$$G = \langle H_1 * H_2; a = \varphi(a)(a \in A_1) \rangle.$$

6.1. Finite (H_1, H_2) -automata

The classical embedding (102) leads naturally to the following adaptations of the definitions from Section 4 to the case of an amalgamated free product.

Let $\mathcal{F}_i \subseteq \mathcal{P}(H_i)$ be a set of subsets of H_i ($i \in \{1, 2\}$) such that every singleton $\{a\}$ with $a \in A_i$ belongs to \mathcal{F}_i . A finite (H_1, H_2) -automaton over $H_1 * H_2$, with labelling sets $(\mathcal{F}_1, \mathcal{F}_2)$ is a 5-tuple

$$\mathcal{A} = \langle \mathcal{L}, \mathcal{Q}, \delta, \mathsf{l}, \mathsf{T} \rangle,$$

where \mathcal{L} is finite and

$$\mathcal{L} \subseteq \mathcal{F}_1 \cup \mathcal{F}_2,$$

\mathcal{Q} is a finite set of states, $\mathsf{l} \subseteq \mathcal{Q}$ is the set of initial states, $\mathsf{T} \subseteq \mathcal{Q}$ is the set of terminal states, and δ (the set of transitions) is a subset of $\mathcal{Q} \times \mathcal{L} \times \mathcal{Q}$ such that for every state $q \in \mathcal{Q}$ there exists $L \in \mathcal{L}$ with $(q, L, q) \in \delta$ and $1 \in L$. For the last point, 1 denotes both the neutral element of H_1 and H_2 . The representation map $\mu_{\mathcal{A}} : H_1 * H_2 \rightarrow \text{BR}(\mathcal{Q})$ is defined as follows. First define for every $h \in H_1 \cup H_2$:

$$\mu_{\mathcal{A},0}(h) = \{(q, r) \in \mathcal{Q} \times \mathcal{Q} \mid \exists L \in \mathcal{L} : (q, L, r) \in \delta \wedge h \in L\}. \quad (103)$$

Now, for every $s \in H_1 * H_2$ of the form (100), let

$$\mu_{\mathcal{A}}(s) = \mu_{\mathcal{A},0}(h_0) \circ \mu_{\mathcal{A},0}(k_1) \circ \mu_{\mathcal{A},0}(h_1) \circ \cdots \circ \mu_{\mathcal{A},0}(k_n) \circ \mu_{\mathcal{A},0}(h_n). \quad (104)$$

The set $L(\mathcal{A}) \subseteq H_1 * H_2$ recognized by \mathcal{A} is defined as for HNN-extensions.

Let us introduce a finite set \mathcal{T}_4 of 4 types, which will induce a partition of the set of states of a finite (H_1, H_2) -automaton:



$$\mathcal{T}_4 = \{(A_1, H_1), (A_2, H_2), (1, H_2), (1, 1)\}. \quad (105)$$

We define a graph $\mathcal{G}_4 = (\mathcal{T}_4, \mathcal{E}_4)$ by:

$$\begin{aligned} \mathcal{E}_4 = & \{((C, D), C, (C, D)) \mid (C, D) \in \mathcal{T}_4\} \\ & \cup \{((A_i, H_i), H_i, (A_{3-i}, H_{3-i})) \mid i \in \{1, 2\}\} \\ & \cup \{((A_2, H_2), H_2, (1, 1)), ((1, H_2), H_2, (A_1, H_1)), ((1, H_2), H_2, (1, 1))\}. \end{aligned}$$

The graph \mathcal{G}_4 is represented in Figure 2. An analogous graph $\mathcal{R}_4 = (\mathcal{T}_4, \mathcal{E}'_4)$ is defined by

$$\begin{aligned} \mathcal{E}'_4 = & \mathcal{E}_4 \setminus \{((A_1, H_1), H_1, (A_2, H_2)), ((A_2, H_2), H_2, (A_1, H_1))\} \\ & \cup \{((A_1, H_1), H_1 \setminus A_1, (A_2, H_2)), ((A_2, H_2), H_2 \setminus A_2, (A_1, H_1))\}, \end{aligned}$$


 Fig. 6. The graph $\mathcal{G}_4 = (\mathcal{T}_4, \mathcal{E}_4)$

 Fig. 7. The graph $\mathcal{R}_4 = (\mathcal{T}_4, \mathcal{E}'_4)$

see Figure 7. The set $\widehat{\mathcal{E}}_4$ is defined analogously to $\widehat{\mathcal{E}}_6$. The reader might be surprised by the asymmetry in the definition of \mathcal{G}_4 and \mathcal{R}_4 : it corresponds to our choice for the definition of an (H_1, H_2) -sequence in (100), where we require $h_0, h_n \in H_2$. This choice was made arbitrarily; we could have also required $h_0, h_n \in H_1$.

A *partitioned finite (H_1, H_2) -automaton* with labelling sets $(\mathcal{F}_1, \mathcal{F}_2)$ is a 6-tuple

$$\mathcal{A} = \langle \mathcal{L}, \mathcal{Q}, \tau, \delta, \mathfrak{l}, \mathfrak{T} \rangle,$$

where \mathcal{L} , \mathcal{Q} , δ , \mathfrak{l} , and \mathfrak{T} are as above and $\tau : \mathcal{Q} \rightarrow \mathcal{T}_4$ assigns a type with every state such that:

$$\begin{aligned} \forall (q, L, r) \in \delta : \{q\} \times L \times \{r\} &\subseteq \widehat{\mathcal{E}}_4 \\ \tau(\mathfrak{l}) = \{(1, H_2)\} &\quad \wedge \quad \tau(\mathfrak{T}) = \{(1, 1)\} \end{aligned}$$

Let us consider the equivalences \sim and \approx over $H_1 * H_2$ defined in Section 5. The notions of \sim -saturated and \approx -compatible finite (H_1, H_2) -automata are defined as in Definition 9, but with the equivalences \approx and \sim being those defined over $H_1 * H_2$.

The notion of a *deterministic* and partitioned finite (H_1, H_2) -automaton \mathcal{A} with labelling sets $(\mathcal{F}_1, \mathcal{F}_2)$ is still defined by the properties (33) and (34). For every $q \in \mathcal{Q}$, and $\Theta \in \mathcal{T}_4$ we again define the set $L^1(q, \Theta) \subseteq H_1 \cup H_2$ by (35). Then, a partitioned finite (H_1, H_2) -automaton \mathcal{A} is *complete* if

$$\forall (\Theta, C, \Theta') \in \mathcal{E}'_4 \quad \forall q \in \tau^{-1}(\Theta) : C \subseteq L^1(q, \Theta').$$

6.2. Normalization of automata

The following three propositions can be now proved in exactly the same way as the corresponding statements for HNN-extensions (Propositions 22, 28 and 33).

Proposition 48. *From a given partitioned and \approx -compatible finite (H_1, H_2) -automaton \mathcal{A} with labelling sets $(\mathcal{F}_1, \mathcal{F}_2)$ one can effectively construct a partitioned, \approx -compatible, \sim -saturated, and unitary (H_1, H_2) -automaton \mathcal{B} with labelling sets $(\{aFa' \mid F \in \mathcal{F}_1, a, a' \in A_1\}, \{aFa' \mid F \in \mathcal{F}_2, a, a' \in A_2\})$ and such that $[L(\mathcal{A})]_{\approx} \cap \text{Red}(H_1, H_2) = L(\mathcal{B}) \cap \text{Red}(H_1, H_2)$.*

Proposition 49. *From a given partitioned, \approx -compatible, and \sim -saturated finite (H_1, H_2) -automaton \mathcal{B} with labelling sets $(\mathcal{F}_1, \mathcal{F}_2)$ one can effectively construct a partitioned, \sim -saturated, deterministic, and complete finite (H_1, H_2) -automaton \mathcal{C} with labelling sets $(\text{Bool}(\mathcal{F}_1), \text{Bool}(\mathcal{F}_2))$ and such that $[L(\mathcal{A})]_{\approx} \cap \text{Red}(H_1, H_2) = L(\mathcal{C})$.*

Proposition 50. *Let $K \subseteq G$. The following are equivalent:*

- (1) K is a rational subset of G .
- (2) $K = \pi(R)$ for some rational subset $R \subseteq (X_1 \cup X_2)^*$.
- (3) $\pi_G^{-1}(K) \cap \text{Red}(H_1, H_2) = L(\mathcal{B}) \cap \text{Red}(H_1, H_2)$ for some partitioned, \approx -compatible, and \sim -saturated finite (H_1, H_2) -automaton \mathcal{B} with labelling sets $(\text{Rat}(H_1), \text{Rat}(H_2))$.

Moreover, in point (3), if the membership problem for languages in $\text{Rat}(H_1)$ and $\text{Rat}(H_2)$, respectively, is decidable, then \mathcal{B} can be computed effectively from a finite automaton for R in point (2).

6.3. Transfer results for rational sets

Using Propositions 48, 49 and 50, the following transfer results for amalgamated free products can be shown in exactly the same way as for HNN-extensions.

Theorem 51. *Let $G = \langle H_1 * H_2; a = \varphi(a)(a \in A_1) \rangle$, where H_1 and H_2 are countably-generated monoids with subgroups $A_1 \leq H_1$ and $A_2 \subseteq H_2$ and $\varphi : A_1 \rightarrow A_2$ is an isomorphism. Then we have:*

- (1) *If $\text{Rat}(H_1)$ and $\text{Rat}(H_2)$ are closed under intersection, then also $\text{Rat}(G)$ is closed under intersection.*
- (2) *If $\text{Rat}(H_1)$ and $\text{Rat}(H_2)$ are closed under complement, then also $\text{Rat}(G)$ is closed under complement.*
- (3) *If the membership problem is decidable for languages in $\text{Rat}(H_1)$ and $\text{Rat}(H_2)$, then the same is true for languages in $\text{Rat}(G)$.*
- (4) *If the emptiness problem is decidable for languages in the boolean closure of $\text{Rat}(H_1)$ and $\text{Rat}(H_2)$, then the same is true for languages in the boolean closure of $\text{Rat}(G)$.*

Statement (3) is covered by [KSS06, Theorem 5.1] (for free products it was shown in [Ned00]) for the case that H_1 and H_2 (and hence G) are groups, whereas (1), (2), and (4) are new to the knowledge of the authors.

6.4. Transfer results for finitely generated subgroups

Let us assume in this section that H_1 and H_2 (and hence G in (99)) are groups. The following subsets $S_1 \subseteq \mathcal{P}(H_1)$ and $S_2 \subseteq \mathcal{P}(H_2)$ serve for labelling finite (H_1, H_2) -automata:

$$\begin{aligned} S_1 &= \{Kh \mid K \text{ is a finitely generated subgroup of } H_1, h \in H_1\} \\ S_2 &= \{Kh \mid K \text{ is a finitely generated subgroup of } H_2, h \in H_2\} \end{aligned}$$

Proposition 52. *Let K be a subgroup of G . The following are equivalent:*

- (1) K is finitely generated.
- (2) K is rational.
- (3) There exists some finite automaton \mathcal{A} over the free monoid $(X_1 \cup X_2)^*$ such that $K = \pi(\mathcal{L}(\mathcal{A}))$.
- (4) $\pi_G^{-1}(K) \cap \text{Red}(H_1, H_2) = \mathcal{L}(\mathcal{B}) \cap \text{Red}(H_1, H_2)$ for some partitioned, \approx -compatible, and \sim -saturated finite (H_1, H_2) -automaton \mathcal{B} with labelling sets (S_1, S_2) .

Moreover, in (4), if the generalized word problem for H_1 and H_2 is decidable, then \mathcal{B} can be constructed effectively from \mathcal{A} in (3).

Theorem 53. *Let $G = \langle H_1 * H_2; a = \varphi(a)(a \in A_1) \rangle$, where H_1 and H_2 are countably-generated groups with subgroups $A_1 \leq H_1$ and $A_2 \leq H_2$ and $\varphi : A_1 \rightarrow A_2$ is an isomorphism. Then we have:*

- (1) If H_1 and H_2 have a decidable generalized word problem, then also G has a decidable generalized word problem.
- (2) If H_1 and H_2 have the Howson property, then also G has the Howson property.
- (3) If H_1 and H_2 have the LERF property, then also G has the LERF property.

Statement (1) is covered by [KWM05, Corollary 5.15], statement (2) was shown in [Bez98,KS70], and statement (3) was shown in [AG73].

Remark 54. There is an alternative way to deduce Theorem 53: using the embedding (102) together with the Theorem 43 on HNN-extensions it suffices to prove that (i) decidability of the generalized word problem, (ii) the Howson property, and (iii) the LERF property are preserved under (i) passing to finitely generated subgroups and (ii) free products. Preservation under passing to finitely generated subgroups is obvious. Preservation under free products can be proved using automata theoretic constructions, which are similar but much simpler than our constructions for amalgamated free products.

For Theorem 51 about rational subsets, the same line of arguments can be applied to the membership problem for rational subsets and the emptiness problem for boolean combinations of rational subsets (decidability of these problems is clearly preserved under passing to finitely generated submonoids). On the other hand, it not clear whether these arguments also apply to the closure under intersection and

complement, respectively, because it is not clear whether these closure properties are preserved under passing to finitely generated submonoids.

Acknowledgments We would like to thank Volker Diekert, Benjamin Steinberg, and the anonymous referee for their useful comments.

References

- [AG73] R.B.J.T. Allenby and R.J. Gregorac. On locally extended residually finite groups. In *Conference on Group Theory (Univ. Wisconsin-Parkside, Kenosha, Wis., 1972)*, number 319 in Lecture Notes in Mathematics, pages 9–17. Springer, 1973.
- [AH89] I.J. Jan Aalbersberg and H.J. Hoogeboom. Characterizations of the decidability of some problems for regular trace languages. *Math. Systems Theory*, 22(1):1–19, 1989.
- [AS75] A.W. Anissimow and F.D. Seifert. Zur algebraischen Charakteristik der durch kontext-freie Sprachen definierten Gruppen. *Elektron. Informationsverarbeitung. Kybernetik*, 11(10–12):695–702, 1975.
- [Ben69] M. Benoist. Parties rationnelles du groupe libre. *C. R. Acad. Sci. Paris, Sér. A*, 269:1188–1190, 1969.
- [Bez98] V.N. Bezverkhniĭ. On the intersection of subgroups in HNN-groups. *Fundam. Prikl. Mat.*, 4(1):199–222, 1998.
- [BJW82] R.V. Book, M. Jantzen, and C. Wrathall. Monadic Thue systems. *Theoret. Comput. Sci.*, 19(3):231–251, 1982.
- [DJ91] N. Dershowitz and J.P. Jouannaud. Rewrite systems. In *Handbook of theoretical computer science*, vol.B, Chapter 2, pages 243–320. Elsevier, 1991.
- [DD90] W. Dicks and M.J. Dunwoody. *Groups acting on graphs*. Cambridge University Press, 1990.
- [Die05] V. Diekert. *Personal communication*. 2005.
- [Git97] R. Gitik. Graphs and separability properties of groups. *J. Algebra*, 188(1):125–143, 1997.
- [Gru90] Z. Grunschlag. *Algorithms in Geometric Group Theory*. PhD thesis, University of California at Berkeley, 1990.
- [Hal49] M. Hall, Jr. Coset representations in free groups. *Trans. Amer. Math. Soc.*, 67:421–432, 1949.
- [HNN49] G. Higman, B.H. Neumann, and H. Neumann. Embedding theorems for groups. *J. London Math. Soc.*, 24:247–254, 1949.
- [How63] J. M. Howie. Embedding theorems for semigroups. *The Quarterly Journal of Mathematics. Oxford. Second Series*, 14:254–258, 1963.
- [KSS06] M. Kambites, P.V. Silva, and B. Steinberg. On the rational subset problem for groups. *J. Algebra*, 309(2):622–639, 2007.
- [KWM05] I. Kapovich, R. Weidmann, and A. Myasnikov. Foldings, graphs of groups and the membership problem. *Internat. J. Algebra Comput.*, 15(1):95–128, 2005.
- [KS70] A. Karrass and D. Solitar. The subgroups of a free product of two groups with an amalgamated subgroup. *Trans. Amer. Math. Soc.*, 150:227–255, 1970.
- [KS71] A. Karrass and D. Solitar. Subgroups of HNN groups and groups with one defining relation. *Canad. J. Math.*, 23:627–643, 1971.
- [LS77] R.C. Lyndon and P.E. Schupp. *Combinatorial group theory*. Springer-Verlag, Berlin, 1977. *Ergebnisse der Mathematik und ihrer Grenzgebiete, Band 89*.

- [LS06a] M. Lohrey and G. Sénizergues. Equations in HNN-extensions. Manuscript, 2006.
- [LS06b] M. Lohrey and G. Sénizergues. Positive theories of HNN-extensions and amalgamated free products. Manuscript, 2006.
- [ME07a] L. Markus-Epstein. Algorithmic problems in amalgams of finite groups. Technical report, arXiv.org, 2007. <http://arxiv.org/abs/0705.0759>.
- [ME07b] L. Markus-Epstein. Stallings' foldings and subgroups of amalgams of finite groups. Technical report, arXiv.org, 2007. <http://arxiv.org/abs/0705.0754>.
- [MR04] V. Metaftsis and E. Raptis. Subgroup separability of graphs of abelian groups. *Proc. Amer. Math. Soc.*, 132(7):1873–1884, 2004.
- [Ned00] M.Yu. Nedbaj. The occurrence problem in a rational subset of the free product of groups. *Vestn. Omsk. Univ.*, 2000(2):17–18, 2000.
- [Reu85] Ch. Reutenauer. Sur les semi-groupes vérifiant le théorème de Kleene. *RAIRO Inform. Théor.*, 19(3):281–291, 1985.
- [Sak87] J. Sakarovitch. On regular trace languages. *Theoret. Comput. Sci.*, 52:59–75, 1987.
- [Sak92] J. Sakarovitch. The “last” decision problem for rational trace languages. In *Proceedings of the 1st Latin American Symposium on Theoretical Informatics (LATIN'92)*, number 583 in Lecture Notes in Computer Science, pages 460–473. Springer, 1992.
- [Sco78] P. Scott. Subgroups of surface groups are almost geometric. *J. London Math. Soc. (2)*, 17(3):555–565, 1978.
- [Sén94] G. Sénizergues. Formal languages and word-rewriting. In *Term Rewriting, Font-Romeu 93, advanced course*, number 909 in Lecture Notes in Computer Science, pages 75–94. Springer, 1994.
- [Sén96] G. Sénizergues. On the rational subsets of the free group. *Acta Inform.*, 33(3):281–296, 1996.
- [Ser03] J.-P. Serre. *Trees*. Springer Monographs in Mathematics. Springer-Verlag, Berlin, 2003. Translated from the French original by J. Stillwell, Corrected 2nd printing of the 1980 English translation.
- [Sta71] J.R. Stallings. *Group Theory and Three-Dimensional Manifolds*. number 4 in Yale Mathematical Monographs. Yale University Press, 1971.
- [Sta83] John R. Stallings. Topology of finite graphs. *Invent. Math.*, 71(3):551–565, 1983.
- [Wis00] D. T. Wise. Subgroup separability of graphs of free groups with cyclic edge groups. *Q. J. Math.*, 51(1):107–129, 2000.

Appendix A. Normal forms

We check here that the classical notion of normal form for HNN-extensions of *groups* adapts to the case of *monoids*, provided that A and B are subgroups of the base monoid. This will result in a proof of Lemma 5.

Let us fix a monoid H , isomorphic subgroups A and B of H , and an isomorphism $\varphi : A \rightarrow B$, as considered in Section 3 and let G be the corresponding HNN-extension defined by (7).

For every subgroup C of H , the left-congruence \equiv_C is defined by

$$h \equiv_C h' \iff \exists c \in C : h = h'c.$$

Since C is a group, \equiv_C is indeed an equivalence relation. The equivalence classes

of \equiv_C are the sets hC for $h \in H$. Let us choose some systems of representatives R_A (resp. R_B) for the equivalence \equiv_A (resp. \equiv_B). We suppose that $1 \in R_A \cap R_B$. Hence, $A \cap R_A = \{1\} = B \cap R_B$. We define now the set of normal forms.

Definition 55. A normal form is an element of $H * \{t, t^{-1}\}^*$ of the form

$$s = h_0 t^{\alpha_1} h_1 \cdots t^{\alpha_i} h_i \cdots t^{\alpha_n} h_n, \quad (\text{A.1})$$

where $n \geq 0$, such that, for every $i \in \{1, \dots, n\}$

- (i) h_n is an arbitrary element of H ,
- (ii) if $\alpha_i = 1$ then $h_{i-1} \in R_A$,
- (iii) if $\alpha_i = -1$ then $h_{i-1} \in R_B$, and
- (iv) if $\alpha_i = -\alpha_{i+1}$ then $h_i \neq 1$.

One can notice that, due to conditions (ii)–(iv), every normal form is a *reduced* sequence. We denote by \mathcal{N} the set of all normal forms. We define a right-action \odot of the monoid $H * \{t, t^{-1}\}^*$ over \mathcal{N} as follows. Let s be a normal form of the form (A.1). For every $h \in H$ we set

$$s \odot h = h_0 t^{\alpha_1} h_1 \cdots t^{\alpha_i} h_i \cdots t^{\alpha_n} (h_n h).$$

The right-action of t on s is defined through 3 cases:

Case (t0). $n = 0$ and $h_0 = r_0 a$ with $r_0 \in R_A$ and $a \in A$:

$$s \odot t = (r_0 a) \odot t = r_0 t \varphi(a)$$

Case (t1). $n > 0, \alpha_n = 1, h_n = r_n a$ with $r_n \in R_A, a \in A$:

$$s \odot t = h_0 t^{\alpha_1} h_1 \cdots t^{\alpha_i} h_i \cdots t^{\alpha_n} r_n t \varphi(a)$$

Case (t2). $n > 0, \alpha_n = -1, h_n = r_n a$ with $r_n \in R_A \setminus \{1\}, a \in A$:

$$s \odot t = h_0 t^{\alpha_1} h_1 \cdots t^{\alpha_i} h_i \cdots t^{\alpha_n} r_n t \varphi(a)$$

Case (t3). $n > 0, \alpha_n = -1, h_n \in A$:

$$s \odot t = h_0 t^{\alpha_1} h_1 \cdots t^{\alpha_i} h_i \cdots t^{\alpha_{n-1}} (h_{n-1} \varphi(h_n))$$

The right-action of t^{-1} on the normal form s is defined similarly through 3 cases:

Case (t⁻¹0). $n = 0$

Case (t⁻¹1). $n > 0, \alpha_n = -1, h_n = r_n b$ with $r_n \in R_B, b \in B$

Case (t⁻¹2). $n > 0, \alpha_n = 1, h_n = r_n b$ with $r_n \in R_B \setminus \{1\}, b \in B$

Case (t⁻¹3). $n > 0, \alpha_n = 1, h_n \in B$

One can easily check that for every $s \in \mathcal{N}$:

$$s \odot (tt^{-1}) = s \odot (t^{-1}t) = s \quad (\text{A.3})$$

Let us compute the sequence $s \odot tbt^{-1}$, when s fulfils case (t1) or (t2):

$$\begin{aligned}
 s \odot tbt^{-1} &= (h_0 t^{\alpha_1} h_1 \cdots t^{\alpha_i} h_i \cdots t^{\alpha_n} r_n t \varphi(a)) \odot (bt^{-1}) \\
 &= (h_0 t^{\alpha_1} h_1 \cdots t^{\alpha_i} h_i \cdots t^{\alpha_n} r_n t(\varphi(a)b)) \odot t^{-1} \\
 &= h_0 t^{\alpha_1} h_1 \cdots t^{\alpha_i} h_i \cdots t^{\alpha_n} (r_n \varphi^{-1}(\varphi(a)b)) \\
 &= h_0 t^{\alpha_1} h_1 \cdots t^{\alpha_i} h_i \cdots t^{\alpha_n} (r_n a \varphi^{-1}(b)) \\
 &= h_0 t^{\alpha_1} h_1 \cdots t^{\alpha_i} h_i \cdots t^{\alpha_n} (h_n \varphi^{-1}(b)) \\
 &= (h_0 t^{\alpha_1} h_1 \cdots t^{\alpha_i} h_i \cdots t^{\alpha_n} h_n) \odot \varphi^{-1}(b).
 \end{aligned}$$

When the sequence s fulfils case (t3) we obtain:

$$\begin{aligned}
 s \odot tbt^{-1} &= (h_0 t^{\alpha_1} h_1 \cdots t^{\alpha_i} h_i \cdots t^{\alpha_{n-1}} (h_{n-1} \varphi(h_n))) \odot (bt^{-1}) \\
 &= (h_0 t^{\alpha_1} h_1 \cdots t^{\alpha_i} h_i \cdots t^{\alpha_{n-1}} (h_{n-1} \varphi(h_n)b)) \odot t^{-1} \\
 &= h_0 t^{\alpha_1} h_1 \cdots t^{\alpha_i} h_i \cdots t^{\alpha_{n-1}} h_{n-1} t^{-1} (h_n \varphi^{-1}(b)) \\
 &= (h_0 t^{\alpha_1} h_1 \cdots t^{\alpha_i} h_i \cdots t^{\alpha_n} h_n) \odot \varphi^{-1}(b)
 \end{aligned}$$

Finally in case (t0) we have

$$\begin{aligned}
 s \odot tbt^{-1} &= (r_0 a) \odot tbt^{-1} = (r_0 t(\varphi(a)b)) \odot t^{-1} = r_0 a \varphi^{-1}(b) \\
 &= s \odot \varphi^{-1}(b).
 \end{aligned}$$

We have thus established that for every $s \in \mathcal{N}$:

$$s \odot (tbt^{-1}) = s \odot \varphi^{-1}(b) \quad (\text{A.4})$$

The set of relations $\{(tt^{-1}, 1), (t^{-1}t, 1)\} \cup \{(tbt^{-1}, \varphi^{-1}(b)) \mid b \in B\}$ generates the congruence \approx over $H * \{t, t^{-1}\}^*$. Equations (A.3) and (A.4) thus show that the action \odot induces a right-action of the monoid G over \mathcal{N} that we still denote by \odot .

Lemma 56. *For every $s \in \mathcal{N}$, $1 \odot \pi_G(s) = s$.*

This lemma can be proved by induction over the integer n in the description (A.1) of a normal form s .

Theorem 57. *The following holds:*

- (1) *For every reduced t -sequence $s \in \text{Red}(H, t)$, there exists $s' \in \mathcal{N}$ with $s \sim s'$.*
- (2) *The set \mathcal{N} is a system of representatives for the congruence \approx over $H * \{t, t^{-1}\}^*$.*

Proof. Let us first prove (1). Let s be some reduced sequence given by (15). Let us show by induction over n that s is \sim -equivalent to some normal form.

If $n = 0$ in (15) then s is itself a normal form. Now assume that $n = m + 1$. By the induction hypothesis, there exist $h'_0, \dots, h'_m \in H$ such that

$$h_0 t^{\alpha_1} h_1 \cdots t^{\alpha_i} h_i \cdots t^{\alpha_m} h_m \sim h'_0 t^{\alpha_1} h'_1 \cdots t^{\alpha_i} h'_i \cdots t^{\alpha_m} h'_m \in \mathcal{N}.$$

First, suppose that $\alpha_n = 1$. Let us decompose $h'_m = r_m a$ with $r_m \in R_A$ and $a \in A$, and set

$$s' = h'_0 t^{\alpha_1} h'_1 \dots t^{\alpha_i} h'_i \dots t^{\alpha_m} r_m t^{\alpha_n} (\varphi(a) \cdot h_n).$$

We have $s \sim s' \in \mathcal{N}$ (note that $t^{\alpha_n} = t \sim at\varphi(a)^{-1}$).

If $\alpha_n = -1$, then we decompose $h'_m = r_m b$ with $r_m \in R_B$ and $b \in B$, and we set

$$s' = h'_0 t^{\alpha_1} h'_1 \dots t^{\alpha_i} h'_i \dots t^{\alpha_m} r_m t^{\alpha_n} (\varphi^{-1}(b) \cdot h_n).$$

Again, we have $s \sim s' \in \mathcal{N}$.

For point (2) in the theorem let us first show that every sequence s is \approx -equivalent to some normal form. As we already saw in Section 3, for every t -sequence s there exists $s' \in \text{Red}(H, t)$ such that $s \approx s'$. By point (1) there exists $s'' \in \mathcal{N}$ with $s'' \sim s'$. Since $\sim \subseteq \approx$, we have $s \approx s'' \in \mathcal{N}$.

It remains to show that two different normal forms are not \approx -equivalent. Let $s, s' \in \mathcal{N}$ and $s \neq s'$. By Lemma 56,

$$1 \odot \pi_G(s) = s, \quad 1 \odot \pi_G(s') = s'.$$

Hence, $\pi_G(s) \neq \pi_G(s')$, i.e., $s \not\approx s'$. \square

We are ready now to give the announced proof of Lemma 5. Suppose that s_1 and s_2 are reduced t -sequences such that $s_1 \approx s_2$. By point (1) of Theorem 57, there exist $s'_1, s'_2 \in \mathcal{N}$ such that $s_i \sim s'_i$ (for $i \in \{1, 2\}$). Since $s'_1 \approx s'_2$, by point (2) of Theorem 57 we have $s'_1 = s'_2$. Finally, $s_1 \sim s'_1 = s'_2 \sim s_2$, which proves that $s_1 \sim s_2$.

Appendix B. Examples of monoids

We give here some examples of monoids \mathbb{M} such that $\text{Rat}(\mathbb{M})$ is closed under intersection but not closed under complement.

Example 58 (Die05). Let $\mathbb{M} = X^*$ be the free monoid over some infinite alphabet X . Since every rational expression uses only a finite number of symbols, every rational set of \mathbb{M} is included in some finitely generated submonoid of \mathbb{M} . Hence, for every $R \in \text{Rat}(\mathbb{M})$, $X^* \setminus R \notin \text{Rat}(\mathbb{M})$.

On the other hand, if R and R' are rational sets of \mathbb{M} , they are rational subsets of some free submonoid Y^* generated by a finite subalphabet $Y \subseteq X$. By Kleene's theorem about rational sets of a finitely generated free monoid, $R \cap R' \in \text{Rat}(Y^*)$, and hence, $R \cap R' \in \text{Rat}(\mathbb{M})$.

Example 59. Using the same arguments as in Example 58 one can show that also for the free group $\mathbb{G} = F(X)$, generated by an infinite set X , $\text{Rat}(\mathbb{G})$ is closed under intersection but $\text{Rat}(\mathbb{G})$ is not closed under complement. One has to use the fact that for a finitely generated free group, the set of rational subsets is a boolean algebra [Ben69].

Let us construct now examples of *finitely generated* monoids.

Example 60. Let $X = \{a, b, c, d, e\}$. Let $L \subseteq \{b, c\}^*$ be some context-free language with non context-free complement. Such examples are abundant, for example one could take

$$L = \{b^p c b^q c b^r \mid p, q, r \in \mathbb{N}, p \neq q \vee q \neq r\}.$$

We then define a semi-Thue system S and a monoid \mathbb{M} by:

$$S = aLd \times \{e\}, \quad \mathbb{M} = X^* / \leftrightarrow_S^*$$

Lemma 61. *The set $\text{Rat}(\mathbb{M})$ is closed under intersection.*

Proof. Let us denote by $\pi : X^* \rightarrow \mathbb{M}$ the canonical projection. Let $R, R' \in \text{Rat}(\mathbb{M})$. There exist $K, K' \in \text{Rat}(X^*)$ such that

$$R = \pi(K) \text{ and } R' = \pi(K').$$

Since the system S has no overlapping left-hand sides nor two rules with the same left-hand side but different right-hand sides, the relation \rightarrow_S^* is confluent (see our recalls in Section 2). It follows that

$$R \cap R' = \pi(K \rightarrow_S^* \cap K' \rightarrow_S^*). \quad (\text{B.1})$$

Since the semi-Thue system S is monadic, both sets of descendants $K \rightarrow_S^*$ and $K' \rightarrow_S^*$ are rational subsets of X^* [BJW82,Sén94]. Equality (B.1) thus shows that $R \cap R'$ is a rational subset of \mathbb{M} . \square

Lemma 62. *For every $R \in \text{Rat}(\mathbb{M})$, the language $\pi^{-1}(R)$ is context-free.*

Proof. Let us assume that $R = \pi(K)$ for some $K \in \text{Rat}(X^*)$. By the confluence property of \rightarrow_S ,

$$\pi^{-1}(R) = \{u \in X^* \mid \exists v \in K \rightarrow_S^* : u \rightarrow_S^* v\}.$$

We already observed that $K \rightarrow_S^*$ is rational. Since the set aLd is context-free, for every rational set Q , $Q \leftarrow_S^*$ is context-free. Applying this property to $Q = K \rightarrow_S^*$ we get that $\pi^{-1}(R)$ is context-free. \square

Lemma 63. *The set $\text{Rat}(\mathbb{M})$ is not closed under complement.*

Proof. Let $R = \{\pi(e)\}$. Clearly, R is a rational subset of \mathbb{M} . Let $R' = \mathbb{M} \setminus R$. One can check that

$$\pi^{-1}(R') \cap a\{b, c\}^*d = a(\{b, c\}^* \setminus L)d.$$

Since $\{b, c\}^* \setminus L$ is not context-free, $\pi^{-1}(R')$ is not context-free (by the closure of context-free languages by intersection with rational sets). By Lemma 62 we can conclude that R' is not rational. \square

We don't know, whether there exists a finitely generated (or even finitely presented) group G such that $\text{Rat}(G)$ is closed under intersection but not closed under complement.